http://www.nap.edu/catalog/11931.html
We ship printed books within 1 business day; personal PDFs are available immediately.

Recognition and Alleviation of Distress in Laboratory Animals

Committee on Recognition and Alleviation of Distress in Laboratory Animals, National Research Council

ISBN: 0-309-10818-7, 132 pages, 6 x 9, (2008)

This PDF is available from the National Academies Press at:
http://www.nap.edu/catalog/11931.html

Visit the National Academies Press online, the authoritative source for all books

and Medicine

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

THE NATIONAL ACADEMIES PRESS 500 Fifth Street, N.W. Washington, DC 20001

NOTICE: The project that is the subject of this report was approved by the Governing Board of the National Research Council, whose members are drawn from the councils of the National Academy of Sciences, the National Academy of Engineering, and the Institute of Medicine. The members of the Committee responsible for the report were chosen for their special competences and with regard for appropriate balance.

This study was supported by the American Veterinary Medical Association; Aventis Pharmaceuticals; the Bosack-Kruger Foundation; Bristol-Myers Squibb; GlaxoSmithKline; Humane Society of the United States; Scientists Center for Animal Welfare; Wyeth-Ayerst Pharmaceuticals; Department of Agriculture, Animal and Plant Health Inspection Service; and Department of Health and Human Services, National Institutes of Health through Contract Number N01-OD-4-2139 Task Order #161. Any opinions, “ndings, conclusions, or recommendations Th24 T0 Tn19.944 0 T6 Tact Number d of

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

COMMITTEE ON RECOGNITION AND ALLEVIATION OF DISTRESS IN LABORATORY ANIMALS

Members

Peter A. Ward (Chair), University of Michigan Medical School
Robert Blanchard, University of Hawaii
Valerie Bolivar, New York State Department of Health
Marilyn J. Brown, Charles River Laboratories
Fon Chang, AstraZeneca, Boston
James P. Herman, University of Cincinnati
Robert Hubrecht, Universities Federation for Animal Welfare
David M. Lawson, Wayne State University School of Medicine (retired)
Steven F. Maier, University of Colorado
David Morton , University of Birmingham, United Kingdom (retired)
Steven M. Niemi, Massachusetts General Hospital
Melinda A. Novak, University of Massachusetts
Stephen L. Zawistowski, American Society for the Prevention of Cruelty to Animals

Staff

Joanne Zurlo, Director >
Lida Anestidou, Study Director (from November 2006) >
Marsha Barrett, Former Project Assistant (to September 2006) >
Kathleen Beil, Administrative Coordinator >
Cameron Fletcher, Senior Editor (from June 2007) >
Rhonda Haycraft, Senior Project Assistant (from September 2006) >
Jennifer Obernier, Former Study Director (to August 2006) >
Susan Vaupel, Former Senior Editor (to June 2007) >
Shimere Williams, Mirzayan Fellow >

Copyright © National Academy of Sciences. All rights reserved.

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

INSTITUTE FOR LABORATORY ANIMAL RESEARCH COUNCIL

Members

Stephen W. Barthold (Chair), University of California, Center for Comparative Medicine, Davis, California
Kathryn A. Bayne, Association for Assessment and Accreditation of Laboratory Animal Care International, Waikoloa, Hawaii
Myrtle A. Davis, Lilly Research Laboratories, Green“eld, Indiana
Jeffrey I. Everitt, GlaxoSmithKline Research and Development, Comparative Medicine and Investigator Support, Research Triangle Park, North Carolina
Michael F. Festing Leicestershire, United Kingdom (to June 2006)
James G. Fox, Massachusetts Institute of Technology, Division of Comparative Medicine, Cambridge, Massachusetts
Nelson L. Garnett, Consultant, Laboratory Animal Care and Use Programs, Dickerson, Maryland
Estelle B. Gauda, Johns Hopkins University School of Medicine, Johns Hopkins Hospital, Baltimore, Maryland
Coenraad F.M. Hendriksen, Netherlands Vaccine Institute, Bilthoven, The Netherlands
Jon H. Kaas, Vanderbilt University, Nashville, Tennessee
Joseph W. Kemnitz, University of Wisconsin, Primate Research Center, Madison, Wisconsin
Judy A. MacArthur Clarkfornia

Guide for the Care and Use of Laboratory Animals„Chinese Version (1996)
Guide for the Care and Use of Laboratory Animals„Spanish Version (1996)
Guide for the Care and Use of Laboratory Animals„Russian Version (1996)
Guide for the Care and Use of Laboratory Animals„French Version (1996)
Guide for the Care and Use of Laboratory Animals„Taiwanese Edition (1996)
Guide for the Care and Use of Laboratory Animals„Portuguese Edition (1996)
Guide for the Care and Use of Laboratory Animals„Japanese Edition (1996)
Rodents (1996)
Nutrient Requirements of Laboratory Animals, Fourth Revised Edition (1995)
Laboratory Animal Management: Dogs (1994)
Recognition and Alleviation of Pain and Distress in Laboratory Animals (1992)
Education and Training in the Care and Use of Laboratory Animals: A Guide for Developing Institutional Programs (1991)
Companion Guide to Infectious Diseases of Mice and Rats (1991)
Infectious Diseases of Mice and Rats (1991)
Immunode“cient Rodents: A Guide to Their Immunobiology, Husbandry, and Use (1989)
Use of Laboratory Animals in Biomedical and Behavioral Research (1988)
Animals for Research: A Directory of Sources, Tenth Edition and Supplement (1979)
Amphibians: Guidelines for the Breeding,Care and Management of Laboratory Animals (1974)

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

Preface >

The impetus for this project was a letter from the New Jersey Associa<

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

Contents >

Summary 1 >
Scope of the Study, 1 >
Stress versus Distress, 2 >
Recognition and Assessment of Stress and Distress, 3 >
Avoiding, Minimizing, and Alleviating Distress, 5 >
Future Studies and Recommendations, 6 >
References, 8 >

1 Introduction 9 >
Regulations Governing Pain and Distress in Laboratory Animals, 9 >
Organization of the Report, 11 >
References, 12 >

2 Stress and Distress: De“nitions 13 >
Introduction, 13 >
What Is Stress?, 14 >
What Is Distress?, 15 >
Implications for Animal Welfare, 17 >
Implications for Research, 21 >
Conclusions, 22 >
References, 22 >

Copyright © National Academy of Sciences. All rights reserved.

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

3 Recognition and Assessment of Stress and Distress 25 >
Introduction, 25 >
Behavioral Recognition of Stress and Distress, 26 >
Physiologic Measures of Stress and Distress, 36 >
Assessment of Distress, 42 >
References, 44 >
Additional References, 52 >

4 Avoiding, Minimizing, and Alleviating Distress 63 >
Introduction, 63 >
Avoiding or Minimizing Distress in Laboratory Animal Care, 64 >
Avoiding or Minimizing Distress in Laboratory Animal Use, 70 >
Alleviating Distress in Laboratory Animals, 75 >
Studying Distress, 79 >
References, 80 >

5 Topics for Further Investigation and Recommendations 87 >
Research Directions, 87 >
Recommendations, 90 >
References, 92 >

Appendix 95 >
Tools to Monitor and Assess Health Status and >
Well-Being in Stress and Distress, 95 >

About the Authors 113 >

Index 117 >iP Tit <</M

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

Summary >

This report is the “rst of two reports prepared as an update to the 1992 National Research Council (NRC) report*Recognition and Alleviation of Pain and Distress in Laboratory Animals*. In the 15 years since the “rst NRC publication on this subject, there has been considerable scienti“c prow1wTh as

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

Both stress and distress are meaningful terms that describe a state of being. While the biological responses to stress are better understood, the scientific, regulatory, and animal welfare communities disagree with respect to a universally accepted definition of distress. Although most definitions of distress characterize it as an aversive, negative state in which coping and adaptation processes in response to stressors fail to return an organism to physiological and/or psychological homeostasis, philosophical differences center on the inclusion of emotions and feelings affected by this state of being. Similarly, while it is accepted that failure of the organism to return to homeostasis adversely impacts an animal's well-being and leads to poor welfare, defining well-being without relying on some form of anthropomorphic measures is a challenge. Scientific research does not yet support objective criteria or principles with which to qualify distress, objective scientific assessment of subjective emotional states cannot be made, and while there is often a measure of agreement on the interpretation of physiologic and/or behavioral variables as indicators of stress, distress, or welfare status, there is not always a direct link. Further, the Committee postulates that even if a universally accepted definition existed, it could not be applied across all species and all conditions, because of the differential impact of the strain, age, gender, genetic background, and environment.

The transition to distress, which occurs when the body cannot cope against the assault of one or more stressors, depends on several factors. Of clear importance are stressor duration, stressor intensity, and the capacity of the individual animal to respond; changes in any of these increases the likelihood of behavioral or physical signs of distress. Thus, minor perturbations may be stressful and/or negatively affect an animal's moment-to-moment emotional state but they would not impair its adaptive capacity and therefore not cause distress (this may be unrelated to the state of the animal's welfare as illustrated in Figure 2-2). In contrast, a major homeostatic disruption (e.g., postsurgical infection), which causes measurable behavioral (e.g., withdrawal) and physiological (e.g., fever) changes that impair an animal's adaptive capacity, would be considered distressful and indicative of poor welfare. However, distress may not manifest itself with recognizable "maladaptive behaviors, such as abnormal feeding or aggression" (NRC 2003a, page 16) but instead begin with subclinical pathological changes (e.g., hypertension or immunosuppression) that can lead to overt disease. These physiological concepts should be integrated within and evaluated in concert with animal welfare principles.

RECOGNITION AND ASSESSMENT OF STRESS AND DISTRESS

While there are some specific behavioral measures of stress, relatively

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

FUTURE STUDIES AND RECOMMENDATIONS

Many questions in the “eld of laboratory animal distress remain unan< swered. The Committee, therefore, offers the following suggestions for research directions that can improve our understanding of distress:

- € @determine whether there arebiomarkers of distress that may be easily measured;
- € @use genomic and proteomic technologies to study the physiology and pathophysiology of stress and distress;
- € @develop possible distress predictorsto be used as outcomes scores (i.e., to predict severity in clinical outcomes, mortality, etc., and adopt humane or surrogate endpoints)or laboratory animals, simi< lar to the predictive severity scoring system used in human inten< sive care units;
- €@ delineate the mechanisms of possible associations between stress/distress and disease behaviorsor abnormal behaviors (e.g., €ste@eotypies);

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

1 >

Introduction >

In 1992, the National Research Council published a report titled

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

Utilization and Care of Vertebrate Animals Used in Testing, Research, and

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

endpoints as well as the use of appropriate statistics and experimental design to minimize the number of animals that must be used. Investigators should establish a partnership with the veterinarians and animal care staff and make provisions in cases of either anticipated or unanticipated distress.

The intent of this report is to assist investigators, veterinarians, animal care staff, and IACUCs in understanding distress so that it can be recognized, alleviated, or prevented. The Committee urges readers to consider the information in this report very carefully and to exercise professional judgment in evaluating situations where distress occurs or is likely to occur. As new information becomes available, it should be incorporated into practice and decision making in the care of laboratory animals. The Committee hopes that this report will be useful in assisting readers to comply with regulations, to achieve reliable scientific outcomes, and, especially, to provide the best possible care for their animal subjects.

REFERENCES

DHHS (Department of Health and Human Services),

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

2 >

Stress and Distress: De“nitions >

INTRODUCTION

The various views and language used in the discussion of stress and distress lead to confusion in the scienti“c, regulatory, and animal welfare communities. Indeed, the animal welfare literature itself does not distin< guish stress from distress in any systematic fashion, and the term distress rarely appears in the biomedical sciences literature. In addition, the U.S.

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

lent phrase is •pain, suffering, distress, and lasting harm.Ž Distress can be

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

animals results in glucocorticoid levels at the diurnal peak that can rival those measured following stressor exposure (Dallman et al. 1987). Thus, no single parameter can serve as a litmus test for stress and diagnosis of stress based on a single metric can be misleading.

- Stressors may not necessarily be unpleasant (defined by the animal's willingness to terminate the stressor); they can be pleasurable (Selye's "eustress" concept; Selye 1975), as defined by an organism's willingness to obtain the stressor. For example, naturally rewarding behaviors, such as exercise, increase sympathetic activity and circulating glucocorticoids in a profile very similar to that seen following aversive stressors (Droste et al. 2003). The self-administration of a drug, such as cocaine, similarly fits the definition of a stressor because multiple physiological systems are recruited to adapt to and oppose the drug's action.
- Physiological and behavioral responses are stressor-specific and so the processes engaged to restore homeostasis or well-being also

1hey celicit alrible vioral rnd ysiologicaresponses : v
0 79>>BDC 1.802 0 Td [(PRsponses atottressorsber 12(3)(12(e)31-7(ei)-(t 2(e)-(e)-(ur)2(1)37(v)2(1)5eig

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html



FIGURE 2-1 Reprinted with permission from Macmillan Publishers Ltd: [Lab Animal] (Moberg 1999), copyright (1999). Prolonged or severe stress depletes bodily reserves and affects normal functions thus requiring extended time to revert to homeostasis. During the recovery period, the animal•s well-being and welfare are compromised and the period of distress will last until the biological resources are suf“ciently replenished (Moberg 2000). The shift in biological resources, such as stunted growth in distressed young animals (Moberg 1999), or evidence of maladaptive behaviors (NRC 1992) that occur in this general scheme of transition to and establishment of distress could be useful in recognizing distress (NRC 2003a, page 21).

proposed in his 1999 paper •When Does Stress Become DistressŽ, the use of reserve resources to cope with prolonged or severe stress has a negative impact on other bodily functions (including behavior) and leads to distress. In the hypothetical scheme depicted in Figure 2-1, the •biological cost of distressŽ requires a prolonged recovery period to revert to homeostasis (Carstens and Moberg 2000).

IMPLICATIONS FOR ANIMAL WELFARE

Current understanding of animal welfare as a measure of the animals• quality of life exists in the context of the social and cultural history of animal care and use as well as an expanding knowledge base related to

Copyright © National Academy of Sciences. All rights reserved.

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

provision of housing without appropriate enrichment).[4] In order to prevent poor or deteriorating welfare, researchers, animal care staff, and institutions have a responsibility to provide high-quality care for laboratory animals, including ready access to fresh water and a nutritive diet; an environment that ensures shelter and comfort; prevention as well as rapid diagnosis and alleviation (as appropriate) of pain, injury, and disease; species-appropriate space, facilities, and (if appropriate) companionship; and conditions and treatment that do not cause negative emotional states. Fraser and colleagues suggest that good animal welfare implies the absence of pain, fear, and hunger; enables a high level of biological functioning (i.e., normal growth, freedom from disease); and (more controversially) enables animals to experience positive emotional experiences such as comfort and contentment (Fraser et al. 1997).

It is possible for an animal to be in a state of poor health that does not impinge on its welfare or emotional state and that may even last for some time without the animal's conscious awareness. For example, an animal might have a life-threatening aneurysm but be unaware of it and therefore not experience a negative emotional state. In the longer term, however, a breakdown in an animal's ability to cope with its environment is likely to lead to adverse emotional states and poor welfare. Some of these cases may be quite minor and not give rise to significant ethical concerns; but prolonged or intense circumstances would compromise the animal's welfare enough to warrant concern and also significantly affect the research results.

An attempt to graphically depict the relationship between distress and welfare is shown in Figure 2-2. Whereas minor perturbations (e.g., short-term restraint of a rodent) affect an animal's welfare in terms of its moment-to-moment emotional state, they do not impair its adaptive capacity and thus do not cause distress. In contrast, a major homeostatic disruption (e.g., postsurgical infection), which causes measurable behavioral (withdrawal) and physiological (fever) changes that impair the adaptive capacity of the animal, is considered "distressful" and is indicative of "poor welfare."

Onset of distress can be difficult to recognize. A safe assumption is to follow the fourth principle of the U.S. Government Principles for Utilization and Care of Vertebrate Animals used in Testing, Research and Teaching: "Proper use of animals, including the avoidance or minimization of discomfort, distress, and pain when consistent with sound scientific practices, is imperative. Unless the contrary is established, investigators should consider that procedures that cause pain or distress in human beings

[4]For more information on the effects of housing on brain function or enrichment see Chapter 3. Additional information is contained in articles by the behaviorists Joseph Garner, Hanno Würbel, and Georgia Mason.

Copyright © National Academy of Sciences. All rights reserved.

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html



FIGURE 2-2 Hypothetical depiction of the relationship of stress, distress, adap< tive capacity, and animal welfare. An animal•squality of life may be progressively deteriorating while it is still successfully coping with a stressor. The precise moment of transition to the maladaptive state or what precipitates it is unknown, but once the tipping point is attained, the deterioration into a severely sick or debilitated animal occurs fairly quickly. At this point, welfare conditions are very poor and immediate ameliorative action is necessary.

may cause pain or distress in other animalsŽ (IRAC 1985). A degree of critical anthropomorphism, outlined above and in the writings of Morton and colleagues (Morton et al. 1990),coupled with behavioral assessments will likely provide the most direct understanding of an animal•s response to a stressor. Useful indicators include the animal•s choice to continue or stop feeding while in a stressful situation, choice tests that demonstrate how (non)aversive a particular stressor is, or demand studies that titrate the extent of the animal•s attraction or aversion to a potential stressor. These gauges of avoidance or aversion may be complemented by physiological data measuring elevated hypothalamic-pituitary-adrenocortical axis (HPA) or sympathoadrenomedullary system (SAM) activity (gene or protein activa< tion), elevated hormone levels, or increased activity in target organs (e.g.,

Copyright © National Academy of Sciences. All rights reserved.

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

heart rate, blood pressure, glucose levels). Many authors have pointed to the desirability of using multiple measures to obtain a more comprehensive data set (Rushen 1991). It is important to underscore that reliance on a single measurement of stress may result in erroneous conclusions. Chapter 3 provides more details on distress recognition.

IMPLICATIONS FOR RESEARCH

There is a rich literature documenting the interference of stress in behavioral and/or physiological endpoints. Strong evidence in rodents has shown that mild stress of 2-3 months duration, a regimen that produces no signs of overt distress, alters the animals' performance in tests of anxiety, depression, and memory (D'Aquila et al. 1994; Rossler et al. 2000; Song et al. 2006; Willner 1997). Other findings indicate that rats' habituation to a test environment can dramatically affect their response to a toxic substance (Damon et al. 1986). On the other hand, in some cases (such as lower anxiety behavior in the elevated plus maze) the effects of stress may actually be beneficial to the experimental procedure, indicating that prolonged stress may not be uniformly detrimental. Chapter 3 documents the contamination of experimental data by unwanted or uncontrolled stress due to inadequate husbandry, noisy environments, olfactory stimuli, or other factors.

The impact of distress on both animal welfare and research results is likely even more pronounced than that of stress. Animals exposed to prolonged severe stress experience underlying changes in physiological functions (e.g., gastric lesions [Ushijima 1985] or immunosuppression [Tournier 2001]) that can interfere with experimental manipulations; alter experimental variables such as behavior (Morton and Griffiths 1985), drug dosing (Saranteas et al. 2004) and clearance; change the progress of a disease (Johnson et al. 2006); and contribute to morbidity and mortality. A variety of stressors can contribute to unintended distress, from postoperative pain or infection to barren housing conditions or the solitary confinement of an individual of a social species (Gunn and Morton 1995; Morton et al. 1993). Stereotypies, abnormal repetitive behaviors indicative of poor well-being (Garner et al. 2003) that are often observed in distressed animals, are thought to reflect defective brain function (Würbel 2001) and to be a result of poor animal welfare (Mason and Latham 2004). Stereotypies are thus likely to interfere with behavioral, neuroscience, and pharmacological studies[5].

The impact of stress and distress on the quality of scientific research can result in the generation of compromised data, which in turn necessitates the use of more animals. This outcome is inconsistent with two of the

[5]It should be noted that the negative connotations of stereotypies are not elevated plus T* y

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

Dallman, M. F., S. F. Akana, C. S. Cascio, D. N. Darlington, L. Jacobson, and N. Levin. 1987. Regulation of ACTH secretion: Variations on a theme of B. Recent Prog Horm Res 43:113-173.

Damon, E. G., A. F. Eidson, C. H. Hobbs, and F. F. Hann. 1986. Effect of acclimation to caging on nephrotoxic response of rats to uranium. Lab Anim Sci 36(1):24-27.

D•Aquila, P. S., P. Brain, and P. Willner. 1994. Effects of chronic mild stress on performance in behavioural tests relevant to anxiety and depression. Physiol Behav 56(5):861-867.

Droste, S. K., A. Gesing, S. Ulbricht, M. B. Muller, A. C. Linthorst, and J. M. Reul. 2003. Effects of long-term voluntary exercise on the mouse hypothalamic-pituitary-adrenocortical axis. Endocrinology 144(7):3012-3023.

Duncan, I. J. H. 2005. Science-based assessment of animal welfare: Farm animals. Rev Sci Tech OIE 24(2):483-492.

EEC (European Economic Commission). 1986. Council Directive 86/609/EEC on the Protection of Animals Used for Experimental and Other Scienti“c Purposes. Available at ec.europa.eu/food/fs/aw/aw_legislation/scienti“c/86-609-eec_en.pdf. Accessed January 30, 2008.

Fraser, D., D. M. Weary, E. A. Pajor, and B. N. Milligan. 1997. A scienti“c conception of animal welfare that re”ects ethical concerns. Anim Welfare 6(3):187-205.

Garner, J. P., C. L. Meehan, and J. A. Mench. 2003. Stereotypies in caged parrots, schizophrenia and autism: Evidence for a common mechanism. Behav Brain Res 145(1-2):125-134.

Gillis S., G. R. Crabtree, and K.A. Smith. 1979. Glucocorticoid-induced inhibition of T cell growth factor production. I. The effect on mitogen-induced lymphocyte proliferation. J Immunol 123(4):1624-1631.

Gunn, D. and D. B. Morton. 1995. Twenty-four hour study of the behavior of New Zealand White rabbits in cages. Appl Anim Behav Sci 45:277-292.

Hedrick, H. and G. Bullock, eds. 2004. The Laboratory Mouse. San Diego: Elsevier Academic Press.

IRAC (Interagency Research Animal Committee). 1985. The U.S. Government Principles for the Utilization and Care of Vertebrate Animals Used in Testing, Research, and Training. Federal Register Vol. 50, No. 97 (May 20, 1985) Of“ce of Science and Technology Policy. Available at: http://grants.nih.gov/grants/olaw/references/phspol.htm#USGovPrinciples. Accessed January 30, 2008.

Johnson, R. R., T. W. Prentice, P. Bridegam, C. R. Young, A. J. Steelman, T. H. Welsh, C. J. Welsh, and M. W. Meagher. 2006. Social stress alters the severity and onset of the chronic phase of Theiler•s virus infection. J Neuroimmunol 175(1-2):39-51.

Kemnitz, J. W., E. B. Roecker, R. Weindruch, D. F. Elson, S. T. Baum, and R. N. Bergman. 1994. Dietary restriction increases insulin sensitivity and lowers blood glucose in rhesus monkeys. Am J Physiol-Endoc M 266(4):E540…E547.

Kirkwood, J. K. 2007. Quality of life: The heart of the matter. Anim Welfare 16:3-7.

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

Moberg, G. P. 1987. Problems in de“ning stress and distress in animals. J. Am Vet Med Assoc 191(10):1207-1211.

Moberg, G. P. 1999. When does stress become distress? Lab Anim 28(4):422-426.

Moberg, G. P. 2000. Biological response tostress: Implications for animal welfare. In The Biology of Animal Stress, G. P. Moberg and J. A. Mench, eds. Wallingford, UK: CAB International. 1-21 pp.

Morton, D. B. and P. H. M. Grif“ths. 1985. Guidelines on the recognition of pain, distress and discomfort in experimental animals and an hypothesis for assessment. Vet Record 116(16):431-436.

Morton, D. B., G. M. Burghardt, and J. A. Smith. 1990. Critical anthropomorphism, animal suffering, and the ecological context. Hastings Cent Rep 20(3):S13-S19.

Morton, D. B., M. Jennings , G. R. Batchelor, D. Bell, L. Birke, K. Davies, J. R. Eveleigh, D. Gunn, M. Heath, B. Howard, P. Koder, J. Phillips, T. Poole, A. W. Sainsbury, G. Sales, D. J. A. Smith, M. Stauffacher, and R. J. Turner. 1993. Re“nements in rabbit husbandry. Lab Anim 27(4):301-329.

J. Holbrook. 1984. Physiological f2attions of glucocorticoids in stress and their relations to pharmacological attions. Endocr Rev 5(1):25-44.

NRC (National Researtio9(Washington DC: National 4.45 0 Td (Co2atil).)Tj 0.0024 Tc 4.258 0

NRC. 2003a. Guidelines for the Care and Use of Mammals in Neuroscience and Behavio Researtio8(h)]TJ /T1_1 1 Tf -0.016 Tw 3.801 0 Td [(.)-7(W)-21ashington, DC: The Natio Nuf“eld Co2atil on Bioethics. 2005.

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

3

Recognition and Assessment of Stress and Distress

INTRODUCTION

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

normalcy, therefore, may or may not serve as clinical indicators of a disease state. Various transgenic and knockout mice that exhibit severe behavioral and physiological phenotypes appear abnormalrelative to their control littermates, but arenormal for their genotype. For example, it is appropri< ate to evaluate Huntington•s disease transgenicmice for signs of stress and distress only relative to their own •normalŽ behavior, taking into account their particular genetic makeup, their abnormal motor patterns, and reduced weight gain (Mangiarini et al. 1996).

BEHAVIORAL RECOGNITION OF STRESS AND DISTRESS

Normal Behavior

Many parameters have an effect on species-speci“c normal behavior and should be taken into consideration when behavioral characteristics are used to determine normalcy or the presenceof stress and distress. Animals exhibit a variety of behavioral changes as part of the normal aging process. Males and females differ in the baselinevalues of many stress markers. Inbred murine strains differ in almost every behavioral, sensory, motor, and physiological trait studied and each inbred strain may respond to stress differently. Similar behavioral differencesin response to stress have been observed in primates. Genetically engineered phenotypes need to be considered when assessing stress anddistress in transgenic and knock< out animals24 Tw 6b05[(dging pr)1a -1EMC /P <<vnìé€ &Êá Þåìa‘£ *-Á é€ /ú îü¢

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

as a function of age, gender, physiological state, and genetic constitution. Third, rearing practices and housing environments may affect expected typical behavior.

1. Age: Many young mammals engage in high levels of social play whereas adults rarely do. Thus, play may be normal for young animals but not necessarily so for adults (Ruppenthal et al. 1974; Vanderschuren et al. 1997). All animals display physiological, behavioral, and cognitive changes as they age. Some of these changes, for example changes in coat/hair appearance and locomotor ability, are overt and easily recognizable. Many laboratory animals display an age-related decline in exploratory activity, which is sometimes correlated to weight gain (as observed, for example, in mice of various strains; Ingram 2000; Ingram et al. 1981). In addition, a number of neurosensory, cardiovascular, endocrine, gastrointestinal, musculoskeletal, and reproductive changes occur with aging, some of which cannot be directly observed in the living animal. Such age-related changes (e.g., hearing and vision deficits) have been documented for a number of murine strains (Hawkins et al. 1985; for more references see Additional References), while cognitive deficits have been reported in aging mice and rats. It is postulated that some of these changes may be gender- and strain-related (Decker et al. 1988; Fischer et al. 1992; Frick et al. 2000). Changes in pain sensitivity and in emotional behavior that may have direct implications for stress and distress have also been reported in

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

less studltTRESS

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

range or, in the case of young animals, gain the appropriate body weight for their age.

Behavioral Signs of Distress

It has been suggested that abnormal behavior, such as stereotypies, is a marker for distress (Dawkins 1990). It remains unclear at this time whether any or all abnormal behaviors qualify as indicators of distress. Several alternative (and largely speculative) hypotheses attempt to explain the occurrence of stereotypic behavior in animals (Mason and Latham 2004; Tiefenbacher et al. 2005). Among these, the stimulation hypothesis suggests that when sensory motor input is low, possibly due to existing (i.e., nonstimulating, poor) housing arrangements, animals engage in stereotypic behavior to self-provide increased sensory-motor input (Sherwin 1998). For example, when cage size constrains normal movements, some animals may respond by developing stereotyped pacing in order to satisfy their need for activity (Draper and Bernstein 1963). The habit hypothesis suggests that although stereotypic behavior may have originally arisen in response to stress or distress, it persists as a habit uncoupled from the situation that originally produced it (Dantzer 1986; Mason 1991). Those who favor the arousal reduction hypothesis suggest that stereotypic behavior may serve to calm the animal and thereby avoid distress (reviewed in Mason 1991). Research shows that in some human and nonhuman primates, even more serious forms of abnormal and self-injurious behavior may function to reduce arousal (Tiefenbacher et al. 2005). The arousal reduction hypothesis is consistent with the view that while an underlying stress or distress state may have initially caused abnormal behavior, eliminating the behavior may be neither desirable nor possible because the stereotypy may sometimes prevent the onset of distress.

Preventing the development of stereotyped behavior by providing the animals with species-specific appropriate environments is obviously desirable and likely to result in improved welfare, especially as enrichment "therapy" may reduce but will not cure the abnormal behavior (van Praag et al. 2000; Wolfer et al. 2004). Although recent studies suggest that stereotypical animals may experience psychological distress due to a putative common mechanism between stereotypy, schizophrenia, and autism, the relationship between stereotypic behavior and distress remains largely unknown and is in need of further study (Garner 2006; Garner and Mason 2002; Garner et al. 2003; Mason 2006).

Copyright © National Academy of Sciences. All rights reserved.

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

Behavioral Signs of Stress

As mentioned in Chapter 2, stress is ubiquitous, it can occur in both pleasurable and aversive situations, and its physiological parameters are well established. Our knowledge of the behavioral correlates of stress, however, is considerably smaller. The behavioral changes observed in a stressed animal (as opposed to a distressed one) may be more subtle and variable, depending on the environmental conditions in which the behavior is being evaluated. In addition to recognizing an animal's normal patterns of behavior, the observer must be well trained and knowledgeable about the normal species-specific behavior in the context of species, strain, gender, and physiological state. Types of behavior commonly explored to investigate the presence of stress include open-field activity, movements in an elevated plus maze, changes in innate behaviors (e.g., movement, grooming, feeding, sexual behavior), defensive behaviors (to external threats), and avoidance/escape (Beck and Luine 2002; for more references see Additional References).

PHYSIOLOGIC MEASURES OF STRESS AND DISTRESS

Endocrinological Parameters

One of the primary endocrinological systems involved in the stress response is the hypothalamic-pituitary-adrenal (HPA) axis, which reacts to stress by releasing glucocorticoids[top] d

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

receptors in the adrenal cortex, resulting in the release of glucocorticoids (cortisol or corticosterone), which mobilize energy stores in response to the perceived stress. When the stressor is removed or otherwise adapted to, glucocorticoids bound to receptors in the hypothalamus and pituitary initiate negative feedback that causes a decrease in the production and release of CRH and ACTH, thus terminating the hormonal response and completing the negative feedback loop (Meaney et al. 1996; Miller and O'Callaghan 2002). It should be noted that the HPA axis affects many hormonal and neural systems and plays a role in modulating the immune system.

Both positive and negative stimuli activate the HPA axis with short- and long-lasting effects. For example, exposure to novel stimuli may elicit exploratory behavior and brief activation of the HPA axis. In contrast, prolonged or repeated stressors, such as social separation of young from their mother, generally elicit strong protest reactions and activation of the HPA axis (Levine 2005; Levine and Mody 2003; Vogt et al. 1980). In the first instance, homeostasis is quickly restored, whereas in the latter case animals may be subjected to chronic HPA changes associated with neuroendocrine stress resistance that persists even after animals are returned to their mother (Parker et al. 2006).

Glucocorticoid levels (usually corticosterone in rodents, cortisol in other species) are used as indicators of the strength and impact of a stressor. Meaningful interpretation of these values, however, presents significant challenges. Glucocorticoids are typically measured in blood serum or plasma but can also be quantified in saliva, urine, feces, and hair (Abelson et al. 2005; for more references see Additional References).

Blood sampling requires venipuncture and possibly other stressful procedures such as handling, transport, capture, restraint, needle stick(s), and sedation. Unless animals are habituated to blood sampling, the method itself can activate the HPA axis thereby confounding assay results. A less stressful sampling method involves measuring glucocorticoid levels in hair. Hair samples are obtained by shaving hair from a particular region (usually the nape of the neck) and then shaving the hair once again after a defined

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

variation (Windle et al. 1998b), making it crucial that samples be collected at the same time each day (e.g., at the nadir of the rhythm). In contrast, urine and feces yield an index of stress reactivity over hours or possibly several days (steady-state samples) and are, therefore, less vulnerable to circadian variation. To date, only hair can provide a chronic index of stress covering a period of several months or more. Assessment of cortisol in hair is presumably unaffected by circadian variation and can be obtained at any time of day.

The above information is relevant for understanding and interpreting what might be revealed about stress and distress by examining the activation of the HPA axis. The two most likely ways to assess distress are to (1) examine differences in basal glucocorticoid levels and identify animals outside a "normal range" or (2) obtain glucocorticoid levels before and after the imposition of a stressor. The first approach is problematic because it assumes that a certain concentration of glucocorticoids indicates distress, although there is no scientific evidence to support this assumption. Moreover, as many sampling methods may themselves activate the stress response, there are no standardized ranges for basal glucocorticoid concentrations. The second approach is also problematic for two reasons: first, the putative relationship between the magnitude of change in glucocorticoid concentrations and distress has not been established; and second, both positive and aversive stimuli activate the HPA axis. Finally, the development of stress or distress is not necessarily associated with activation of the HPA axis, as hormonal changes are not necessarily present under all clearly stressful conditions. For example, animals that experience chronic neuropathic pain do not exhibit changes in circadian corticosteroid levels or oscillations in HPA responsivity to restraint, despite the presence of neuropathic pain markers (mechanical allodynia and hyperalgesia) and activation of central pain and stress circuits in the amygdala (Bomholt et al. 2005; Ulrich-Lai et al. 2006).

Other "Stress" Hormones

As is the case with the hormones of the HPA axis, stressors alter the secretion of other endocrine factors (e.g., prolactin, growth hormone, luteinizing hormone, A-melanocyte stimulating hormone [A-MSH], and oxytocin). Serum levels of these hormones can be effectively used to monitor the temporal dynamics of stress responses. While some (prolactin, A-MSH, oxytocin) increase during stress, others decrease (growth hormone, luteinizing hormone, prolactin), depending on the animal species and the physiological state in which stress occurs (Armario et al. 1984; for more references see Additional References). Due to the fact that these hormones

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

dian or circadian variations), it is necessary to take into consideration and control for their normal patterns of secretion in order to accurately interpret their concentration levels. Moreover, their usefulness is subjected to the same limitations as discussed above, although chronic indwelling vascular catheters and automated blood collection systems may circumvent this limitation to some degree (Abelson et al. 2005; for more references see Additional References).

Neurological Parameters

Stressors activate the autonomic nervous system, specific brain areas, and various neurotransmitters, yet a cause and effect relationship has not yet been firmly established. Candidates for the role of a master integrator include the region of the amygdala and the neuropeptide corticotrophin-releasing factor.

The Autonomic Nervous System

Many different types of stressors cause the rapid activation of the sympathetic division of the autonomic nervous system (ANS) (Blanc et al. 1991; for more references see Additional References). This activation leads to increased cardiac output via increased heart rate and stroke volume; redistribution of blood flow from splanchnic, renal, and cutaneous vascular beds to active muscle; increased mobilization of nutrients; and increased heat production. Some stressors may also increase the activity of the parasympathetic division, affecting both core body temperature and the gastrointestinal system (e.g., disturbed intestinal absorption, gastric ulceration, colitis; Johnson et al. 2002; for more references see Additional References).

Direct monitoring of autonomic activity to assess the presence of distress in conscious animals is technically challenging, while indirect measures are somewhat easier to acquire (Lee et al. 1997; Randall et al. 1994; Zhang and Thoren 1998). For example, telemetry in conscious, unrestrained animals is an effective method for the continuous monitoring of physiologic alterations in heart rate, respiration, blood pressure, ECG, and body temperature (Akutsu et al. 2002; for more references see Additional References). Once again, however, changes in these parameters do not necessarily indicate stress as they may result from nonstressful stimuli (e.g., circadian variations).

Neurotransmitters

Considerable effort has been directed at exploring the neurotransmitter systems and brain areas activated in response to stress as different insults

Copyright © National Academy of Sciences. All rights reserved.

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

activate separate but speci“c patterns of transmitters, modulators, and brain locations. An active area of investigation has been the identi“cation of a basic core neural circuit that is activated by all stressors. A common approach has been to divide stressors into categories, hypothesizing that each category would activate a particular set of neural structures. The stressors are classi“ed as either •processiveŽ (i.e., stressors that require interpretation by higher brain structures) or •systemicŽ (i.e., stressors evoked by

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

factors alter the activity in structures that project to the top of the central nucleus of the amygdala (Sotres-Bayon et al. 2006), underscoring the amygdala as the key integrative site for fear.

In contrast, much remains to be learned about the neural control of defense responses to threat. For example, depending on circumstances, external threats such as the presence of a predator may result in ”ight, freez< 5eru, in.0023vISTRESS

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

Assessment of distress varies in relation to the species, husbandry conditions, and experimental protocol employed as well as with each individual animal, and is most effectively achieved by the collection of multiple behavioral and physiological parameters and the use of a team approach that includes researchers, veterinarians, and animal caretakers/technicians. While the most reliable distress measures are the clinical signs previously described, identification and interpretation of these results depends on a solid foundation of knowledge of animal behavior and may likely require special training of relevant personnel.

Distress evaluation becomes crucial in two contexts: (1) when the research protocol calls for the animals' exposure to stressful situations known to produce distress; and (2) when any animal unexpectedly shows signs of distress. In the first case, the experimental protocol approved by the Institutional Animal Care and Use Committee (IACUC) generally includes procedures and decision algorithms for distress management. The appropriate intervention will be informed by the stressor's duration and intensity as well as some of the animal's individual characteristics (e.g., species, age, gender). In the second case, additional assessments and monitoring may be necessary.

Once an animal has shown initial signs of distress, there should be immediate communication between the primary investigator, the veterinarian, and the animal care staff to determine whether the distress is related to the study (whether anticipated or unanticipated) or further investigation into its cause is required. The discussion should also include potential interventions (see Chapter 4) and their effects on the objectives of the research project, as they may introduce unknown variables into the study. Options may include removal of the animal from the study population or euthanasia, depending on the severity and prognosis of the distress insult. It is essential to maintain a collaborative relationship and dialogue between those responsible for the care and welfare of the animal throughout the assessment.

The next step is to identify the etiology or trigger of the distress episode by performing a thorough examination of the animal and its environment. The investigation should begin by obtaining information regarding the animal's species, strain, age, gender, and reproductive status. An effective examination should account for species-related differences among natural behaviors, learning abilities, and levels of intelligence, in addition to the ways animals use their 2(al05s-l(y intro7 T37(a)29Bhe)]TJ 0.0 Td [(T)ee120l0hh)-

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

is an important part of the process, as background information and history may enable the veterinarian to determine whether preexisting medical conditions were resolved. An examination of colony records and interviews with animal care staff may help pinpoint possible environmental triggers. Other causes for consideration include husbandry and handling procedures, the behavior of other animals in the room, temperature variances, noises, vibrations, and odors, as well as any speci“c research-related (i.e., protocol speci“cations) or investigator-related (i.e., disturbance of housing routine) activities.

Clinical signs should initially be examined in a relatively undisturbed animal in order to assess the animal•s natural unprovoked behavior (e.g., appearance, behavior, posture, respiratory rate and pattern). The animal showing signs of distress is then observed more closely followed by gentle handling and examination to measure body weight, body condition and ybehavior i/y weight0.112pattern, w

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

Coe, C. L., G. R. Lubach, W. B. Ershler, and R. G. Klopp. 1989. In"uence of early rearing on lymphocyte proliferation response s in juvenile rhesus monkeys. Brain Behav Immun 3(1):47-60.

Crawley, J. N. 2000. What•s Wrong with My Mouse? Behavioral Phenotyping of Transgenic and Knockout Mice. New York: Wiley-Liss, John Wiley & Sons, Inc.

Crawley, J. N., J. K. Belknap, A. Collins, J. C. Crabbe, W. Frankel, N. Henderson, R. J. Hitzemann, S. C. Maxson, L. L. Miner, A. J. Silva, J. M. Wehner, A. Wynshaw-Boris, and R. Paylor. 1997. Behavioral phenotypes of inbred mouse strains: Implications and recom< mendations for molecular studies. Psychopharmacology 132(2):107-124.

Dalla, C., K. Antoniou, G. Drossopoulou, M. Xagoraris, N. Kokras, A. S"kakis, and Z. Papadopoulou-Daifoti. 2005. Chronic mild stress impact: Are females more vulnerable? Neuroscience 135(3):703-714.

Dantzer, R. 1986. Behavioral, physiological, and functional aspects of stereotyped behavior: A review and reinterpretation. J Anim Sci 62(6):1776-1786.

Dantzer, R. 2004. Cytokine-induced sickness behaviour: A neuroimmune response to activa< tion of innate immunity. Euro J Pharmacol 500(1-3):399-411.

Davenport, M. D., S. Tiefenbacher, C. K. Lutz, M. A. Novak, and J. S. Meyer. 2006. Analysis of endogenous cortisol concentrations in the hair of rhesus macaques. Gen Comp Endocr 147(3):255-261.

Davis, M. and P. J. Whalen. 2001. The amygdala: Vigilance and emotion. Mol Psychiat 6(1):13-24.

Dawkins, M. S. 1990. From an animal•s point of view: Motivation, "tness, and animal welfare. Behav Brain Sci 13(1):1-61.

Deak, T., K. T. Nguyen, M. Fleshner, L. R. Watkins, and S. F. Maier. 1999. Acute stress may facilitate recovery from a subcutaneous bacterial challenge. Neuroimmunomodulation 6(5):344-354.

Decker, M. W., M. A. Pelleymounter, and M. Gallagher. 1988. Effects of training on a spatial memory task on high af"nity choline uptake in hippocampus and cortex in young adult and aged rats. J Neurosci 8(1):90-99.

Dhabar, F. S. and K. Viswanathan. 2005. Short-term stress experienced at time of immuniza< tion induces a long lasting increase in immunological memory. Am J Physiol Reg-I 289: R738-R744.

Draper, W. A. and I. S. Bernstein. 1963. Stereotyped behavior and cage size. Percept Motor Skill 16:231-234.

Ducottet, C. and C. Belzung. 2005. Correlations between behaviours in the elevated plus-maze and sensitivity to unpredictable subchronic mild stress: Evidence from inbred strains of mice. Behav Brain Res 156(1):153-162.

Felten, D. L., S. Y. Felten, S. L. Carlson, J. A. Olschowka, and S. Livnat. 1985. Noradrenergic and peptidergic innervation of lymphoid tissue. J Immunol 135(2 Suppl):755-765.

Figueiredo, H. F., C. M. Dolgas, and J. P. Herman. 2002. Stress activation of cortex and hippo- campus is modulated by sex and stage of estrus. Endocrinology 143(7):2534-2540.

Fischer, W., K. S. Chen, F. H. Gage, and A. Bjorklund. 1992. Progressive decline in spatial learning and integrity of forebrain cholinergic neurons in rats during aging. Neurobiol Aging 13(1):9-23.

Fish, E. W., D. Shahrokh, R. Bagot, C. Caldji, T. Bredy, M. Szyf, and M. J. Meaney. 2004. Epigenetic programming of stress responses through variations in maternal care. Ann NY Acad Sci 1036:167-180.

Fleshner, M., J. Nermann, L. Lockwood, M. L. Laudenslager, L. R. Watkins, and S. F. Maier. 1995. Stressed rats fail to expand the CD45RC+CD4+ (th1-like) T cell subset in response to KLH: Possible involvement of interferon gamma. Brain Behav Immun 9(2):101-112.

Copyright © National Academy of Sciences. All rights reserved.

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

Moy, S. S., J. J. Nadler, A. Perez, R. P. Barbaro, J. M. Johns, T. R. Magnuson, J. Piven, and J. N. Crawley. 2004. Sociability and preference for social novelty in “ve inbred strains: An approach to assess autistic-like behavior in mice. Genes Brain Behav 3(5):287-302.

Muglia, L., L. Jacobson, P. Dikkes, and J. A. Majzoub. 1995. Corticotropin-releasing hor< mone de“ciency reveals major fetal but not adult glucocorticoid need. Nature 373(6513):427-432.

Nash, L. T., J. Fritz, P. A. Alford, and L. Brent. 1999. Variables in”uencing the origins of diverse abnormal behaviors in a large sample of captive chimpanzees (Pan troglodytes). Am J Primatol 48(1):15-29.

Nemeroff, C.B. and W. W. Vale. 2005. The neurobiology of depression: Inroads to treatment and new drug discovery. J Clin Psychiat 66(Suppl 7):5-13.

Novak, M. A. 2003. Self-injurious behavior in rhesus monkeys: New insights into its etiology, physiology, and treatment. Am J Primatol 59(1):3-19.

Novak, M. A. and S. J. Suomi. 1988. Psychological well-being of primates in captivity. Am Psychol 43(10):765-73.

O•Connor, K. A., J. D. Johnson, M. K. Hansen, J. L. Wiesler-Frank, L. R. Watkins, and S. F. Maier. 2003. Peripheral and central proin”ammatory cytokine response to a severe acute stressor. Brain Res 991:123-132.

Olsson, I. A. S. and C. M. Sherwin. 2006. Behaviour of laboratory mice in different housing conditions when allowed to self-administer an anxiolytic. Lab Anim 40(4):392-399.

Parker, K. J., C. L. Buckmaster, K. Sundlass, M. A. Schatzberg, and D. M. Lyons. 2006. Maternal mediation, stress inoculation, and the development of neuroendocrine stress resistance in primates. Proc Nat Acad Sci USA 103(8):3000-3005.

Pezze, M. A. and J. Feldon. 2004. Mesolimbic dopaminergic pathways in fear conditioning. Prog Neurobiol 74(5):301-320.

Randall, D. C., D. R. Brown, L. V. Brown, and J. M. Kilgore. 1994. Sympathetic nervous activity and arterial blood pressure control in conscious rat during rest and behavioral stress. Am J Physiol 267(5 Part 2):R1241-R1249.

Ribeiro, S. C., S. E. Kennedy, Y. R. Smith, C. S. Stohler, and J. K. Zubieta. 2005. Interface of physical and emotional stress regulation through the endogenous opioid system and mu-opioid receptors. Prog Neuropsychopharmacol Biol Psychiatry 29(8):1264-1280.

Ruppenthal, G. C., C. G. Walker, and G. P. Sackett. 1991. Rearing infant monkeys (Macaca nemestrina) in pairs produces de“cient social development compared with rearing in single cages. Am J Primatol 25(2):103-113.

Ruppenthal, G. C., M. K. Harlow, C. D. Eisele, H. F. Harlow, and S. J. Suomi. 1974. Develop< ment of peer interactions of monkeys reared in a nuclear-family environment. Child Dev 45(3):670-682.

Sackett, G. P., G. C. Ruppenthal, and A. E. Davis. 2002. Survival, growth, health, and repro< duction following nursery rearing compared with mother rearing in pigtailed monkeys (Macaca nemestrina). Am J Primatol 56(3):165-183.

Sanders, V. M., D. J. Kasprowicz, A. P. Kohm, and M. A. Swanson. 2001. Neurotransmitter receptors on lymphocytes and other lymphoid cells. In Psychoneuroimmunology, 3rd ed. R. Ader, D. L. Felten, and N. Cohen, eds. New York: Academic Press. 161-197 pp.

Serrats, J. and P. E. Sawchenko. 2006. CNS activational responses to staphylococcal enterotoxin B: T-lymphocyte-dependent immune challenge effects on stress-related circuitry. J. Comp Neurol 495(2):236-254.

Sherwin, C. 1998. Voluntary wheel running: A review and novel interpretation. Anim Behav 56(1):11-27.

Silver, L. M. 1995. Mouse Genetics Concepts and Applications New York: Oxford University Press.

Copyright © National Academy of Sciences. All rights reserved.

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

Smith, L. K. and G. A. Metz. 2005. Dietaryrestriction alters “ne motor function in rats. Physiol Behav 85(5):581-592.

Snowdon, C. T. and A. Savage. 1989Psychological well-being of captive primates: General considerations and examples from callitrichids. In Housing, Care and Psychological Wellbeing of Captive and Laboratory Primates, E. F. Segal, ed. Park Ridge, NJ: Noyes Publications. 75-88 pp.

Sotres-Bayon, F., C. F. Cain, and J. E. LeDoux. 2006. Brain mechanisms of fear extinc< tion: Historical perspectives on the contributions of prefrontal cortex. Biol Psychiat 60(4):329-336.

Stenzel-Poore, M., S. C. Heinrichs, S. Rivest, G. F. Koob, and W. W. Vale. 1994. Over< production of corticotrophin-releasing factor in transgenic mice: A genetic model of anxiogenic behavior. J Neurosci 14(5 Pt 1):2579-2584.

Suomi, S. J. 2004. How gene-environment interactions shape biobehavioral development: Lessons from studies with rhesus monkeys. Res Hum 1:205-222.

Thomas, G. 2002. The S6 kinase signaling pathway in the control of development and growth. Biol Res 35(2):305-313.

Tiefenbacher, S. T., M. A. Novak, C. K.Lutz, and J. S. Meyer. 2005. The physiology and neurochemistry of self-injurious behavior: A nonhuman primate model. Front Biosci 10(1):1-11.

Tournier, J. N., J. Mathieu, Y. Mailfert, E. Multon, C. Drouet, A. Jouan, and E. Drouet. 2001. Chronic restraint stress induces severe disruption of the T-cell speci“c response to tetanus toxin vaccine. Immunology 102(1):515-523.

Ulrich-Lai, Y. M, W. Xie, J. T. Meij, C. M. Dolgas, L. Yu, and J. P. Herman. 2006. Limbic and HPA axis function in an animal model of chronic neuropathic pain. Physiol Behav 88(1-2):67-76.

van Bogaert, M. J., L. Groenink, R. S. Oosting, K. Westphal, G. van der Sluis, T. de Gr< Gr52enink, L. van der Sluis, G. Westphal, K. Oosting, R. S. van Bogaert, M. J. de Gr< 615025 T17

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

Zhang, W. and P. Thoren. 1998. Hyper-responsiveness of adrenal sympathetic nerve activity in spontaneously hypertensive rats to ganglionic blockade, mental stress and neuroglucopenia. Pflugers Arch 437(1):56-60.

ADDITIONAL REFERENCES

Page 26 The identification of species-typical behavior often comes from ethograms developed by researchers to describe the kinds of behavior that animals display in various settings (Bronson 1979).

Chopra, P. K., P. K. Seth, and S. Seth. 1992. Behavioural profile of free-ranging rhesus monkeys. Primate Rep 32:75-105.

Latham, N. and G. Mason. 2004. From house mouse to mouse house: The behavioural biology of free-living Mus musculus and its implications in the laboratory. Appl Anim Behav 86(3-4):261-289.

Mayer, J. 2007. Use of behavior analysis to recognize pain in small animals. Lab Anim 36(6):43-48.

Morton, D. B. 2002. Behaviour of rabbits and rodents. In The Ethology of Domestic Animals, An Introductory Text, Per Jensen, ed. Oxford: CABI Wallingford Oxford. 193-209 pp.

Morton, D. B. and P. H. M. Griffiths. 1985. Guidelines on the recognition of pain, distress and discomfort in experimental animals and an hypothesis for assessment. Vet Record 116(16):431-436.

Olivier B. and D. van Dalen. 1982. Social behaviour in rats and mice: An ethologically based model for differentiating psychoactive drugs. Aggress Behav 8(2):163-168.

Page 27 Such age-related changes (e.g., hearing and vision deficits) have been documented for a number of murine strains (Hawkins et al. 1985).

Johnson, K. R., Q. Y. Zheng, and K. Noben-T

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

Garner, J. P. 2005. Stereotypies and other abnormal repetitive behaviors: Potential impact on validity, reliability, and replicability of scienti“c outcomes. ILAR J 46(2):106-117.
Hubrecht, R. C., J. A. Serpell, and T. B. Poole. 1992. Correlates of pen size and housing condi< tions on the behaviour of kennelled dogs. Appl Anim Behav Sci 34:365-383.
Luescher, U. A., D. B. McKeown, and J. Halip. 1991. Stereotypic or obsessive-compulsive disorders in dogs and cats. Vet Clin North Am-Small 21(2):401-413.
Morton, D. B., M. Jennings, G. R. Batchelor, D. Bell, L. Birke, K. Davies, J. R. Eveleigh, D. Gunn, M. Heath, B. Howard, P. Koder, J. Phillips, T. Poole, A. W. Sainsbury, G. Sales, D. J. A. Smith, M. Stauffacher, and R. J. Turner. 1993. Re“nements in rabbit husbandry. Lab Anim 27(4):301-329.
Waiblinger, E. and B. Konig. 2004. Re“nement of gerbil housing and husbandry in the labora< tory. Anim Welf 13(Suppl):S229-S235.

Page 34 Such protocols are widely used in gerontology research where diet has been shown to slow aging, extend lifespan, and reduce the incidence of age-related diseases in rodents (Goto et al. 2002).

Goto, S., R. Takahashi, Z. Radak, and R. Sharma. 2007. Bene“cial biochemical outcomes of late-onset dietary restriction in rodents. Ann N Y Acad Sci 1100:431-441.
Jamieson, H. A., S. N. Hilmer, V. C. Cogger, A. Warren, R. Cheluvappa, D. R. Abernethy, A. V. Everitt, R. Fraser, R. de Cabo, and D. G. Le Couteur. 2007. Caloric restriction reduces age-related pseudocapillarization of the hepatic sinusoid. Exp Gerontol 42(4):374-378.
Hyun, D. H., S. S. Emerson, D. G. Jo, M. P. Mattson, and R. de Cabo. 2006. Calorie restriction up-regulates the plasma membrane redox system in brain cells and suppresses oxidative stress during aging. Proc Natl Acad Sci USA 103(52):19908-19912.
Seymour, E. M., R. V. Parikh, A. A. Singer, and S. F. Bolling. 2006. Moderate calorie restric< tion improves cardiac remodeling and diastolic dysfunction in the Dahl-SS rat. J Mol Cell Cardiol 41(4):661-668.

Page 36 Types of behavior commonly explored to investigate the presence of stress include open-“eld activity, movements in an elevated plus maze, changes in innate behaviors (e.g., movement, grooming, feeding, sexual behavior), defensive behaviors (to external threats), and avoidance/escape (Beck and Luine 2002).

Blanchard, R. J., C. R. McKittrick, and D. C. Blanchard. 2001. Animal models of social stress: > Effects on behavior and brain neurochemical systems. Physiol Behav 73(3):261-271. >
Ely, D. R., V. Dapper, J. Marasca, J. B. Correa, G. D. Gamaro, M. H. Xavier, M. B. Michalowski, > D. Catelli, R. Rosat, M. B. Ferreira, and C. Dalmaz. 1997. Effect of restraint stress on feeding behavior of rats. Physiol Behav 61(3):395-398.
Harkin, A., T. J. Connor, J. M. O•Donnell, and J. P. Kelly. 2002. Physiological and behavioral responses to stress: What does a rat “nd stressful? Lab Anim 31(4):42-50.
Mangiavacchi, S., F. Masi, S. Scheggi, B. Leggio, M. G. De Montis, and C. Gambarana. 2001. Long-term behavioral and neurochemical effects of chronic stress exposure in rats. J Neurochem 79(6):1113-1121.
Pare, W. P. and S. M. Tejani-Butt. 1996. Effect of stress on the behavior and 5-HT system in Sprague-Dawley and Wistar Kyoto rat strains. Integr Physiol Beh Sci 31(2):112-121.

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

Guo, A. L., F. Petraglia, M. Criscuolo, G. Ficarra, R. E. Nappi, M. Palumbo, A. Valentini, and A. R. Genazzani. 1994. Acute stress- or lipopolysaccharide-induced corticosterone secretion in female rats is independent of the oestrous cycle. Eur J Endocrinol 131(5):535-539.

Hall, F. S., J. M. Sundstrom, J. Lerner, and A. Pert. 2001. Enhanced corticosterone release after a modi“ed forced swim test in Fawn hooded rats is independent of rearing experience. Pharmacol Biochem Be 69(3-4):629-634.

Harper, J. M. and S. N. Austad. 2000. Fecal glucocorticoids: A noninvasive method of measur< ing adrenal activity in wild and captive rodents. Physiol Biochem Zool 73(1):12-22.

Ishikawa, M., S. Ohdo, H. Watanabe, C. Hara, and N. Ogawa. 1995. Alteration in circadian rhythm of plasma corticosterone in rats following sociopsychological stress induced by communication box. Physiol Behav 57(1):41-47.

Kirschbaum, C. and D. H. Hellhammer. 1994. Salivary cortisol in psychoneuroendocrine research: Recent developments and applications. Psychoneuroendocrinology 19(4):313-333.

Klosterman, L. L, J. T. Murai, and P. K. Siiteri. 1986. Cortisol levels, binding, and prop< erties of corticosteroid-binding globulin in the serum of primates. Endocrinology 118(1):424-434.

Ling, S. and F. Jamali. 2003. Effect of cannulation surgery and restraint stress on the plasma corticosterone concentration in the rat: Application of an improved corticosterone HPLC assay. J Pharm Pharm Sci 6(2):246-251.

Livezey, G. T., J. M. Miller, and W.

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

Tuli, J., J. A. Smith, and D. B. Morton. 1995a. Corticosterone, adrenal and spleen weight in mice after tail bleeding, and its effect on nearby animals. Lab Anim 29(1):90-95.

Tuli, J., J. A. Smith, and D. B. Morton. 1995b. Effects of acute and chronic restraint on the adrenal gland weight and serum corticosterone concentration of mice and their faecal output of oocysts after infection with Eimeria apionoides. Res Vet Sci 59(1):82-86.

Tuli, J., J. A. Smith, and D. B. Morton. 1995c. Stress measurements in mice after transportation. Lab Anim 29(2):132-138. seru.

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

Seggie, J. A. and G. M. Brown. 1975. Stress response patterns of plasma corticosterone, prolactin, and growth hormone in the rat, following handling or exposure to novel environment. Can J Physiol Pharmacol 53(4):629-637.

Siegel, R. A., N. Conforti, and I. Chowers. 1980. Neural pathways mediating the prolactin secretory response to acute neurogenic stress in the male rat. Brain Res 198(1):43-53.

Simms, D. D., L. V. Swanson, and R. Bogart. 1978. Effect of collection stress on serum growth hormone levels in pygmy goats. J Anim Sci 46(2):458-462.

Yelvington, D. B., G. K. Weiss, and A. Ratner. 1984. Effect of corticosterone on the prolactin response to psychological and physical stress in rats. Life Sci 35(16):1705-1711.

Page 39 Moreover, their usefulness is subjected to the same limitations as discussed above, although chronic indwelling vascular catheters and automated blood collection systems may circumvent this limitation to some degree (Abelson et al. 2005).

Atkinson, H. C., S. A. Wood, Y. M. Kershaw, E. Bate, and S. L. Lightman. 2006. Diurnal variation in the responsiveness of the hypothalamic-pituitary-adrenal axis of the male rat to noise stress. J Neuroendocrinol 18(7):526-533.

Lightman, S. L., R. J. Windle, M. D. Julian, M.S. Harbuz, N. Shanks, S. A. Wood, Y. M. Kershaw, and C. D. Ingram. 2000. Signi“cance of pulsatility in the HPA axis. Novar Fnd Symp 227:244-257.

Rogers, W. R., J. H. Lucas, B. C. Mikiten, H. D. Smith, and J. L. Orr. 1995. Chronically indwelling venous cannula and automatic blood sampling system for use with nonhuman primates exposed to 60 Hz electric and magnetic “elds. Bioelectromagnetics Suppl 3:103-110.

Windle, R. J., S. A. Wood, N. Shanks, P. Perks, G. L. Conde, A. P. da Costa, C. D. Ingram, and S. L. Lightman. 1997. Endocrine and behavioural responses to noise stress: Comparison of virgin and lactating female rats during non-disrupted mate T1_,lactaG6e3.25 Td (of)18 Neuroendoc

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

Stiedl, O., M. Meyer, T. Kishimoto, M. G. Rosenfeld, and J. Spiess. 2003. Stress-mediated heart rate dynamics after deletion of the gene encoding corticotropin-releasing factor receptor 2. Eur J Neurosci 17(10):2231-2235.

Tornatzky, W. and K. A. Miczek. 1994. Behavioral and autonomic responses to intermittent social stress: Differential protection by clonidine and metoprolol. Psychopharmacology 116(2):346-356.

van den Buuse, M., G. Lambert, M. Fluttert, and N. Eikelis. 2001a. Cardiovascular and behavioural responses to psychological stress in spontaneously hypertensive rats: Effect of treatment with DSP-4. Behav Brain Res 119(2):131-142.

Wood, S. K., R. E. Verhoeven, AZ. Savit, K. C. Rice, P. S.

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

4

Avoiding, Minimizing, and Alleviating Distress

The simplest approach to avoiding, minimizing, and alleviating distress in laboratory animal care and use is to follow the principles of the Three Rs—refinement, reduction, and replacement. It is important, however, to strike a balance between the integrity of research outcomes and the welfare of animals used. Investigators, veterinarians, and animal care personnel should function as a team and base their decisions on professional judgment, best practices, and thorough clinical evaluation of distressed animals. Refining aspects of housing, husbandry, enrichment, and socialization helps alleviate or prevent distress. Refining the experimental design, utilizing humane or surrogate endpoints, and using the appropriate statistical analyses (including an accurate calculation of sample size) helps reduce the numbers of animals used and alleviate some of their distress. A team approach is necessary to address treatment options for distressed animals. When using procedures that intentionally result in distress, the investigator should, in consultation with the veterinarian and the IACUC, develop a plan that will establish limits to the levels of distress allowed.

INTRODUCTION

Established ethical, regulatory, and scientific practices, standards, and policies mandate avoiding animal distress whenever possible. However, if research or regulatory testing objectives cause a research animal to experience distress, it is incumbent upon the animal user to identify the cause(s) of distress, attempt to minimize its duration and intensity, and/or provide the

Copyright © National Academy of Sciences. All rights reserved.

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

means for the animal to cope. The simplest approach is to follow the principles developed by Russell and Burch known as the Three Rs (Russell and Burch 1959). The Three Rs are (1) *refinement* of the protocol to minimize or eradicate distress for the species used (e.g., by employing nonclinical [e.g., molecular measurements] or defined [e.g., tumor growth instead of survival] endpoints; giving positive rewards; changing or refining the data/sample collection methods; or instituting species-specific husbandry refinements such as enrichment); 2) *reduction* of the number of animals used to the absolute minimum necessary (based on appropriate statistical sample size determination or other field-specific methods), particularly if they are likely to experience unavoidable distress; and (3) *replacement* of an animal with a nonanimal model or a less sentient species, usually of a lower phylogenetic order, such as a primitive invertebrate.

As discussed in Chapters 2 and 3, distress may result from a single intense or prolonged stressful experience or from several simultaneous stressors that might individually cause stress but not likely distress. Therefore, mitigating some potentially stressful circumstances, such as husbandry schedules, may allow an animal to better adapt to other stressors such as experimental procedures. It is important to weigh any possibly adverse impact of replacement, refinement, and reduction on scientific outcome against both the negative impact of failure to avoid, minimize, or alleviate stress and distress on the research data and the numbers of potentially wasted animal lives.

This chapter identifies approaches to avoid or minimize distress through the alleviation and minimization of stress, in both the care and use of laboratory animals. The chapter also suggests ways to alleviate distress that cannot be avoided or minimized because of scientifically justified research protocols, and addresses the challenges and compromises that arise when the object of research itself is the study of stress and distress. This chapter is not intended to be comprehensive but to provide investigators and IACUC members with an awareness of common problems and useful strategies. The Committee encourages all who are involved in laboratory animal care and use to think creatively when considering solutions to specific circumstances.

AVOIDING OR MINIMIZING DISTRESS IN LABORATORY ANIMAL CARE

Most animals are able to cope with a relatively wide range of naturally occurring environments, but such environments are not usually characteristic of laboratory animal facilities. Research environments are generally designed as a compromise between the needs of the animal, the user, and

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

the husbandry staff. While most animals continue to function normally[1] in animal care facilities, some do not adapt successfully to a laboratory environment (see Chapter 3) and it is important to consider the effects of

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

Socialization

It is generally appropriate to house naturally gregarious animals in compatible social groups unless there are scientific or welfare reasons not to do so (National Health and Medical Research Council 2004; Canadian Council on Animal Care 1993; Council of Europe 2006; NRC 1996). Social housing can activate stress responses involving the HPA axis in rats, but when a wider range of measures is taken into account, overall, social housing is neither stressful nor harmful (Hurst et al. 1997, 1998). For example, even macaques fitted with a cranial implant could be paired with another compatible macaque without it inflicting damage to the device or interfering with the research goals (Roberts and Platt 2005). Furthermore, a considerable body of evidence indicates that housing naturally sociable animals (e.g., rats, mice, dogs, primates) in solitary conditions can result in stress and harm (Baker 1996; Eaton et al. 1994; Hetts 1991; Hubrecht 1995; Novak 2003; Patterson-Kane 2002; Sharp and Lawson 2003; Van Loo et al. 2004). Even cats, which are not particularly gregarious, can benefit from social housing (Council of Europe 2006). It is therefore important to provide thorough scientific rationale for solitary housing.

Disruption of established social groups, pairing (for additional information see Appendix), or the introduction of animals to larger preformed units are all potential causes of aggression or stress. As a husbandry refinement, therefore, social groups should be established early, and disruption of established groups should be minimal, as demonstrated in studies of mice, rabbits, and cats (Bradshaw and Hall 1999; Jennings et al. 1998; Morton et al. 1993; Sharp et al. 2002b). Close cooperation with the supplier or breeder may be necessary to promote group formation and ensure minimal disruption of group dynamics. Adequate socialization to both humans and conspecifics at an early age may also help prevent subsequent stress and distress (Council of Europe 2006).

Animals housed in social groups generally need adequate space as C3]blishe

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

and other steroids in samples from saliva (Aardal and Holm 1995; Kiess et al. 1995) and plucked hair (Davenport et al. 2006). Even less handling is involved when samples are taken from voided urine, feces, expired air, and shed hair (Poon and Chu 1999), if these methods are validated for the species under study. Other noninvasive techniques for data collection include sound recordings (Holy and Guo 2005), cameras (Hobbs et al. 1997), or noninvasive, sensor-laden apparel simply worn by the animal (Jarrell et al. 2005).

Humane Endpoints

Validated endpoints that occur earlier in the course of the protocol and involve no detectable indication of disease, injury, or abnormal behavior can prevent or minimize distress in experimentation and testing. The use of humane endpoints (i.e., "end a study earlier to avoid or terminate unrelieved pain and/or distress"; Stokes 2000) or surrogate endpoints (i.e., those that can reliably substitute for more distressing or painful phenomena) is especially applicable in scientific disciplines that focus primarily on molecular and cellular phenomena associated with disease (Morton 2000; Hendriksen and Morton 1999). In these cases, biochemical changes may be detectable at early stages in the disease process, prior to the manifestation of clinical signs consistent with distress. For example, elevated white blood cell counts are detectable in leukemia models before illness becomes obvious and serum biochemical values often change in early stages of toxicity before animals appear ill (Poon and Chu 1999). Thus, taking measurements or collecting samples from animals before the appearance of any clinical signs (including all clinical manifestations, not only those related to distress) is desirable, especially when the signs themselves are not the study's focus. In such cases, the predictive value of validated endpoints may permit early euthanasia of these animals and postmortem collection of data or samples (for additional information see Appendix). Alternatively, a clinically normal animal could be anesthetized before a distressful procedure and euthanized before regaining consciousness.

Familiarity with certain procedures or experimental protocols often allows for predicting the course of adverse clinical signs and distress. In many instances death results from indirect effects such as dehydration and is not related to the response variable under study. In mice, for example, progressive hypothermia due to low food intake will cause an animal's death over several days. However, distress can be minimized through the use of validated humane endpoints, such as euthanizing the animals at the first recording of low body temperature (Morton 1998; Soothill et al. 1992). The choice and use of endpoints should be part of the experimental protocol whenever possible.

Copyright © National Academy of Sciences. All rights reserved.

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

a refinement approach that makes it possible to measure tumors so precisely that the animal may be euthanized before any clinical signs arise. This method by itself may greatly reduce the numbers of animals estimated by the sample size determination. Other animal models that benefit from new imaging technologies include those for cardiovascular diseases (labeled cells; Jaffer et al. 2006) and inflammatory bowel diseases (colonoscopy; Becker et al. 2005).

ALLEVIATING DISTRESS IN LABORATORY ANIMALS

As has been noted in Chapter 3, even with reasonable steps to avoid or minimize housing and husbandry-related stressors, distress may still unexpectedly appear once a protocol begins or following a change in husbandry. Many of the steps involved in the alleviation of distress, such as a team management approach and prompt veterinary action, are identical to the procedures described in Chapter 3 for recognizing and assessing the presence of distress. However, before implementing any response plan, the principal investigator/study director and veterinarian or designee should review the objectives of the protocol to determine if the alleviation of distress would adversely affect the research project. Identification of a refinement after approval of a protocol should include amendment of the protocol to adopt this change. If the distress is anticipated or results from a significant husbandry error, regulations require notification of the Institutional Animal Care and Use Committee (IACUC) and possibly of regulatory agencies as well, especially if animal distress results in protocol suspension (DHHS 2005; USDA 2005). Table 4-1 provides an algorithm for responding to unexpected animal distress.

Medical conditions unrelated to the study objectives (e.g., spontaneous self-injurious behavior, fight-related injuries, newly diagnosed ectoparasite infestations) may be treatable without compromising the study. However, additional diagnostic tests may be necessary and even, depending on the therapeutic interventions selected, the removal of the animal from the study, either temporarily or permanently. In addition to eliminating the underlying cause, treatment modalities to address the behavioral signs may include changing the environmental parameters (such as cagemate, caging type, or housing location [Fontenot et al. 2006]; administering analgesics or anxiolytics; engaging in behavior modification and training [Reinhardt 2003; Schapiro et al. 2001]); providing environmental enrichment; dispensing psychotropic medications; or, in severe cases, euthanizing the animal. In one case, environmental enrichment decreased abnormal behaviors in pigtail macaques that could not be socially housed (Kessel and Brent 1998). In contrast, the presence of puzzle feeders, which encouraged manipulation, did not reduce self-injurious behaviors in rhesus monkeys (Novak et

Copyright © National Academy of Sciences. All rights reserved.

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

TABLE 4-1 Example of a decision and response algorithm for unanticipated distress in laboratory animals

6 c^ b Va >hh j Zh	Egd\ gV b >hh j Zh
Egd b eian Xd b b j c^XViZ ^c^i^Va dWhZgkVi^dch id eg^cX^eVa ^ckZhi^\ Vidg$hi j Yn Y^gZXidg! Xa^c^XVa kZiZg^cVg^Vc! [VX^a^in b VcV\ Zg#	Egd b eian VcY VXX j gViZan YdX j b Zci Xa^c^XVa h^\ ch VcY igZVi b Zcih ^c i]Z Vc^ b Va»h gZXdgY#
6 hhZhh Vc^ b Va»h Xa^c^XVa hiVi j h VcY igZVi b Zci dei^dch l ^i] gZheZXi id i]Z egdidXda#	: kVa j ViZ di]Zg Vc^ b Vah dc i]Z hV b Z egdidXda dg]d j hZY cZVgWn id YZiZg b ^cZ ^[b dgZ Vc^ b Vah VgZ edhh^Wan ^c h^ b ^a Vg Y^higZhh#
6 Y b ^c^hiZg Z b Zg\ ZcXn kZiZg^cVgn XVgZ ^[^cY^XViZY VcY V[iZg Xdch j aiVi^dc l ^i] i]Z eg^cX^eVa ^ckZhi^\ Vidg$hi j Yn Y^gZXidg#	9 ZiZg b ^cZ ^[i]Z Y^higZhh VcY VXXd b eVcn^c\ Xa^c^XVa h^\ ch VgZ V XdchZf j ZcXZ d[ZmeZg^ b ZciVi^dc!] j hWVcYgn Zggdg! dg di]Zg XV j hZ#
>[i]Z Vc^ b Va»h XdcY^i^dc ^h \ gVkZ VcY i]Z eg^cX^eVa ^ckZhi^\ Vidg$hi j Yn Y^gZXidg dg YZh^\ cZZ XVccdi WZ XdciVXiZY! i]Z Vc^ b Va b Vn WZ Z j i]Vc^oZY Vi i]Z Y^gZXi^dc d[i]Z Xa^c^XVa kZiZg^cVg^Vc#	G ZY j XZ dg Za^ b ^cViZ i]Z hd j gXZ h d[Y^higZhh! ^[`cd l c VcY ^[Xd b eVi^WaZ l ^i] i]Z V^ b h d[i]Z egdidXda#
>chi^i j iZ egZXV j i^dcVgn b ZVh j gZh VcY h j eedgi^kZ XVgZ ^[^cY^XViZY#	Cdi^[n i]Z >6 8 J 8 VcY edhh^Wan gZ\ ja Vidgn V\ ZcX^Zh d[h^\ c^[^XVci Vc^ b Va Y^higZhh#
	6 b ZcY i]Z egdidXda id Vkd^Y dg gZY j XZ Y^higZhh ^c b dgZ Vc^ b Vah# >[VaiZg^c\ i]Z egdidXda l ^aa Xd b egd b ^hZ hX^Zci^[^X V^ b h dg gZ\ ja Vidgn ZcYed^cih! Vhh^\ c Vc^ b Vah id V b dgZ hZkZgZ eV^c$Y^higZhh XViZ\ dgn#

al. 1998), while Coleman and colleagues demonstrated that the ability of individual monkeys to respond to conventional training methods is closely correlated with their unique temperament (exploratory or inhibited person< alities; Coleman et al. 2005).

The actual causes of distress may also lead to sequelae that require attention even if the underlying cause is not treatable. For example, the clinical signs of a distressed animal often include dehydration and weight loss resulting from anorexia. Provision of supplemental "uids and nutrition may relieve the compounding impact of dehydration or poor body condi< tion on the compromised animal. Supplemental heat, cooling, bedding, social housing, and human companionship are other strategies that make a

Copyright © National Academy of Sciences. All rights reserved.

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

STUDYING DISTRESS

Distress in humans may be more widespread (or at least more readily recognized) than that observed in nonhuman animals because of unique human cognitive capacities, such as the ability to clearly communicate threatening, dangerous, or painful conditions; to remember these circumstances and their consequences over extended periods of time; and to apply the emotions engendered to other stimuli on the basis of verbal or categorical concepts (Sapolsky 1994). A substantial proportion of the American population will at some point suffer from an illness that is distressing or even incapacitating (e.g., depression or a severe anxiety disorder). Many of

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

REFERENCES

Aardal, E. and A. C. Holm. 1995. Cortisol in saliva„reference ranges and relation to cortisol in serum. Eur J Clin Chem Clin 33(12):927-932.

Abeyesinghe, S. M., C. M. Wathes, C. J. Nicol, and J. M. Randall. 2001. The aversion of broiler chickens to concurrent vibrational and thermal stressors. Appl Anim Behav Sci 73(3):199-215.

AVMA (American Veterinary Medical Association). 2007. 2007 AVMA Guidelines on Euthanasia.

Baker, K. C. 1996. Chimpanzees in single cages and small social groups: Effects of housing on behavior. Contemp Top Lab Anim 35(3):71-74.

Bardo, M. T., J. E. Klebaur, J. M. Valone, and C. Deaton. 2001. Environmental enrich< ment decreases intravenous self-administration of amphetamine in female and male rats. Psychopharmacology 155(3):278-284.

Baumans, V. 2005. Environmental enrichment for laboratory rodents and rabbits: Requirements of rodents, rabbits, and research. ILAR J 46(2):162-170.

Bayne, K. 2005. Potential for unintended consequences of environmental enrichment for laboratory animals and research results. ILAR J 46(2):129-139.

Becker, C., M. C. Fantini, S. Wirtz, A. Nikolaev, R. Kiesslich, H. A. Lehr, P. R. Galle, and M. F. Neurath. 2005. In vivo imaging of colitis and colon cancer development in mice using high resolution chromoendoscopy. Gut 54(7):950-954.

Belz, E. E., J. S. Kennell, R. K Czambel, R. T. Rubin, and M. E. Rhodes. 2003. Environmental enrichment lowers stress-responsive hormones in singly housed male and female rats. Pharmacol Biochem Be 76(3-4):481-486.

Benaroya-Milshtein, N., N. Hollander, A. Apter, T. Kukulansky, N. Raz, A. Wilf, I. Yaniv, and C. G. Pick. 2004. Environmental enrichment in mice decreases anxiety, attenuates stress responses and enhances natural killer cell activity. Eur J Neurosci 20(5):1341-1347.

Bene“el, A. C., W. K. Dong, and W. T. Greenough. 2005. Mandatory •enriched Ž housing of laboratory animals: The need for evidence-based evaluation. ILAR J 46(2):95-105.

Blanchard, D. C. and R. J. Blanchard. 2005. Stress and aggressive behaviors. In Biology of Aggression, R. J. Nelson, ed. Oxford: Oxford University Press.

Bradshaw, J. W. S. and S. L. Hall. 1999. Af“liative behaviour of related and unrelated pairs of cats in catteries: A preliminary report. App Anim Behav Sci 63(3):251-255.

Broadbear, J. H., G. Winger, J. E. Rie r, and J. H. Woods. 2004. Corticotropin-releasing hormone antagonists, astressin B and antalarmin: Differing pro“les of activity in rhesus monkeys. Neuropsychopharmacology 29(6):1112-1121.

Canadian Council on Animal Care (CCAC). 1993. Guide to the Care and Use of Experimental Animals, 2nd Edition.

Chung, K. C., L. K. Kalliainen, S. V. Spilson, M. R. Walters, and H. M. Kim. 2002. The preva< lence of negative studies with inadequate statistical power: An analysis of the plastic surgery literature. Plast Reconstr Surg 109(1):1-6.

Clark, J. M. 1997. Biocontainment facilities: Implications for animal care and welfare. In Animal Alternatives, Welfare, and Ethics, L. F. M. van Zutphen and M. Balls, eds. New York: Elsevier.

Clark, V. A. 2002. The prevalence of negative studies with inadequate statistical power: An analysis of the plastic surgery literature by Kevin C. Chung, MD, MS, Loree K. Kalliainen, MD, Sandra V. Spilson, MPH, Madonna R. Walters, MS, RN, Hyungjin Myra Kim, ScD. Plast Reconstr Surg 109(1):7-8.

Clough, G. 1982. Environmental effects on animals used in biomedical research. Biol Rev 57(Pt 3):487-523.

Copyright © National Academy of Sciences. All rights reserved.

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

AVOIDING, MINIMIZING, AND ALLEVIATING DISTRESS

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

Jaffer, F. A., P. Libby, and R. Weissleder. 2006. Molecular and cellular imaging of atherosclerosis: Emerging applications. J Am Coll Cardiol 47(7):1328-1338.

Jarrell, D. M., J. Camacho, D. Funk-Flavin, and S. M. Niemi. 2005. A new method for comprehensive, non-invasive physiological data recording in conscious macaques. Contemp Top Lab Anim 44(4):69.

Jennings, M., G. R. Batchelor, P. F. Brain, A. Dick, H. Elliott, R. J. Francis, R. C. Hubrecht, J. L. Hurst, D. B. Morton, A. G. Peters, R. Raymond, G. D. Sales, C. M. Sherwin, and C. West. 1998. Refining rodent husbandry: The mouse. Lab Anim 32(3):233-259.

Kaldi, I., M. Dittmar, P. Pierce, and R. E. Anderson. 2003. L-NAME protects against acute light damage in albino rats, but not against retinal degeneration in P23H and S334ter transgenic rats. Exp Eye Res 76(4):453-461.

Kessel, A. L. and L. Brent. 1998. Cage toys reduce abnormal behavior in individually housed pigtail macaques. J Appl Anim Welf Sci 1(3):227-234.

pile micemacaques. 43(2Td 281-2924227-234. 22cc1988macaques. at34. MeijJaffp. K.(22ccSommJ)11(af

dac34.

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

Tomei, F., M. G. Ruf“no, E. Tomao, T. P. Baccolo, M. V. Rosati, and F. Strollo. 2000. Acute experimental exposure to noise and hormonal modi“cations. J Environ Sci Heal A 35:537-555.

Tsai, P. P., U. Pachowsky, H. D. Stelzer, and H. Hackbarth. 2002. Impact of environmental enrichment in mice: Effect of housing conditions on body weight, organ weights and haematology in different strains. Lab Anim 36(4):411-419.

Turner, P. V., K. L. Smiler, M. Hargaden, and M. A. Koch. 2003. Re“nements in the care and use of animals in toxicology studies„regulation, validation, and progress. Contemp Top Lab Anim 42(6):8-15.

Vahl, T. P., Y. M. Ulrich-Lai, M. M. Ostrander, C. M. Dolgas, E. E. Elfers, R. J. Seeley, D. A. D•Alessio, and J. P. Herman. 2005. Comparative analysis of ACTH and corticosterone sampling methods in rats. Am J Physiol Endoc M 289(5):E823-828.

USDA (U.S. Department of Agriculture). 2005. 9CFR2.31 Code of Federal Regulations, title 9, Volume 1, part 2, subpart C, section 2.31 (d) (1) (i): Institutional Animal Care and Use Committee. Available at: http://www.aphis.usda.gov/animal_welfare/downloads/awr/9cfr2.31.txt. Accessed January 30, 2008.

Van Loo, P. L., E. Van der Meer, C. L. J. L. Kruitwagen, J. M. Koolhaas, L. F. Van Zutphen, and V. Baumans. 2004. Long-term effects of husbandry procedures on stress-related parameters in male mice of two strains. Lab Anim 38(2):169-177.

Van Loo, P. L., H. A. Van de Weerd, L. F. Van Zutphen, and V. Baumans. 2004. Preference for social contact versus environmental enrichment in male laboratory mice. Lab Anim 38(2):178-188.

Wasowicz, M., C. Morice, P. Ferrari, J. Callebert, and C. Versaux-Botteri. 2002. Long-term effects of light damage on the retina of albino and pigmented rats. Invest Ophth Vis Sci 43(3):813-820.

Waterton, J. C., B. J. Middleton, R. Pickford, CP. Allott, D. Checkley, R. A. Keith. 2000. Reduced animal use in ef“cacy testing in disease models by the use of sequential experimental designs. In Progress in the Reduction, Re“nement and Replacement of Animal Experimentation, M. Balls, A.-M. van Zeller, and M. Halder, eds. Oxford: Elsevier Science.

Weed, J. L. and J. M. Raber. 2005. Balancing animal research with animal well-being: Establishment of goals and harmonization of approaches. ILAR J 46(2):118-128.

Weissleder, R. 2006. Molecular imaging in cancer. Science 312(5777):1168-1171.

Weld, K. P., J. A. Mench, R. A. Woodward, M. S. Bolesta, S. J. Suomi, and J. D. Higley. 1998. Effect of tryptophan treatment on self-biting and central nervous system serotonin metabolism in rhesus monkeys (Macaca mulatta). Neuropsychopharmacology 19(4):314-321.

Würbel, H. 2000. Behaviour and the standardization fallacy. Nat Genet 26(3):263.

Copyright © National Academy of Sciences. All rights reserved.

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

ing them passively, even though the latter approach might result in missing something serious. While there exists a growing body of scienti“c evidence (for example see Bayne et al. 2002; EEC 1986; Kaliste 2004; Morton 2002), it is important to approach continuing attempts to establish what isin the best interests of animals with rigorous scienti“c interdisciplinary methods. Even experts such as veterinarians, ethologists, and animal welfare sci< entists have to guard against the twin traps of anthropomorphism

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

3. @Institutional support for and embrace of a commitment to animal welfare of the laboratory animals is essential. Veterinarians and animal care personnel who work with research animals on a daily basis should have adequate time and contact with the animals to properly evaluate their well-being. Funding for training programs is crucial to the training and development of specialized laboratory animal veterinarians and animal behaviorists and should increase, because in addition to such objective measurements as weight loss or lack of grooming, clinical judgment is vital to effective assess< ments of stress and distress.

4. @Appropriate statistical methodologies are an essential tool for the avoidance, minimization, and alleviation of distress.

5. @There should be a clearinghouse (or some other venue such as a website or a specialized peer-reviewed journal) for publication of research on the effects of enrichment strategies on parameters such as physiology, distress, and endpoints for all laboratory animals (one useful example is the Primate Enrichment Database hosted by the Animal Welfare Institute)[1]. Although a variety of journals (such as Lab Animal, Applied Animal Behaviour Science, Animal Welfare, Laboratory Animals, Contemporary Topics in Lab= oratory Animal Science, Comparative Medicine) publish research pertaining to animal welfare, the highly specialized nature of the "eld makes it dif"cult for the larger scienti"c community to remain informed about recent advances and ongoing debates. Biomedical research journals should be more open to submissions from sci< entists whose research focuses on animal welfare issues so that concerns about research interference or unjusti"ed expenses can be debated on scienti"c, ethical, or regulatory grounds.

6. @Obtaining funding for welfare research is often dif"cult, especially when project applications compete against other "elds of science due to lack of an appropriate/separate research oversight body. In the United Kingdom the funds available for welfare research have increased dramatically with the founding of the National Center for the Replacement, Re"nement and Reduction of Animals in Research (NC3Rs)[2]. In the United States, the National Institutes of Health, Environmental Protection Agency, and other federal insti< tutions have occasionally provided funding to develop or validate

[1]http://www.awionline.org/SearchResultsSite/enrich.aspx. >
[2]NC3Rs website: www.nc3rs.org.uk. >

Copyright © National Academy of Sciences. All rights reserved.

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

nonanimal or nonvertebrate alternatives. Funding for laboratory animal welfare research, however, is usually available only in small amounts from nongovernmental organizations such as the Animal Welfare Institute, the Johns Hopkins Center for Alternatives to Animal Testing, the American College of Laboratory Animal Medicine, and the American Association for Laboratory Animal Science. Given the impact of better animal welfare on science as well as the growing public interest in the treatment of laboratory animals, federal agencies and large foundations that support biomedical and behavioral research should make funds available specifically for the avenues of investigation listed above and for other related topics.

7. Animal welfare scientists and researchers and scientists who use animal models should communicate with each other more frequently in order to compare objectives and progress and to identify opportunities for collaboration. Neutral groups and/or other established research and science policy entities can provide platforms and venues for such exchanges.

REFERENCES

AVMA (American Veterinary Medical Association). 2007. 2007 AVMA Guidelines on Euthanasia.

Bayne, K. A. L., J. A. Mench, B. V. Beaver, and D. B. Morton. 2002. Laboratory Animal Behavior. In: Laboratory Animal Science, ACLAM series 2nd Ed. New York: Academic Press. 1240-1264 pp.

Bradshaw, W. S. and R. A. Casey. 2007. Anthropomorphism and anthropocentrism as influences in the quality of life of companion animals. Anim Welf 16 (suppl): in press.

Brown, M., L. Carbone, K. M. Conlee, M. S. Dawkins, I. J. Duncan, D. Fraser, G. Griffin, V. A. Hampshire, L. A. Lambert, J. A. Mench, D. Morton, J. Richmond, B. E. Rollin, A. N. Rowan, M. L. Stephens, and H. Würbel. 2006. Report of the Working Group on Animal Distress in the Laboratory. Lab Animal 35(8):26-30.

EEC (European Economic Commission). 1986. Council Directive 86/609/EEC on the Protection of Animals Used for Experimental and Other Scientific Purposes. Available at: ec.europa.eu/food/fs/aw/aw_legislation/scientific/86-609-eec_en.pdf. Accessed August 2007.

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

Kaliste, E., ed. 2004. The Welfare of Laboratory Animals. Dordrecht/London: Kluwer Academic Publishers.

Knaus, W. A., J. E. Zimmerman, D. P. Wagner, E. A. Draper, and D. E. Lawrence. 1985. APACHE II—A severity of disease classification system. Crit Care Med 13:818-829.

Knaus, W. A., D. P. Wagner, E. A. Draper, J. E. Zimmerman, M. Bergner, P. G. Bastos, C. A. Sirio, D. J. Murphy, T. Lotring, A. Damiano, et al. 1991. The APACHE III prognostic system. Risk prediction of hospital mortality for critically ill hospitalized adults. Chest 100(6):1619-1636.

McGreevy, P. D., P. J. Cripps, N. P. French, L. E. Green, and C. J. Nicol. 1995. Management factors associated with stereotypic and redirected behavior in the thoroughbred horse. Equine Vet J 27(2):86-91.

Morton, D. B. 2002. Behaviour of rabbits and rodents. In The Ethology of Domestic Animals—An Introductory Text, P. Jensen, ed. Oxford: CABI Wallingford. 193-209 pp.

Nicol, C. J., C. Potzsch, K. Lewis, and L. E. Green. 2003. Matched concurrent case-control study of risk factors for feather pecking in hens on free-range commercial farms in the UK. Br Poultry Sci 44(4):515-523.

Niemi, S. M. 2006. Making comparative medicine more so. J Am Vet Med Assoc 228(2):199-200.

Weisstaub, N. V., M. Zhou, A. Lira, E. Lambe, J. Gonzalez-Maeso, J. P. Hornung, E. Sibille, M. Underwood, S. Itohara, W. T. Dauer, M. S. Ansorge, E. Morelli, J. J. Mann, M. Toth, G. HT00.

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

are introduced by gender, age, physiological state, genetics, and genetic modi“cation of the animals.

The “rst three tables below contain behavioral categories and descrip< tions of physiologic activities in which rhesus macaques, common marmosets, and rabbits engage. In order to determine what kind of behavior it is that an animal exhibits, one needs to be knowledgeable in the ethology and hus< bandry of the species in question. For example, aggression may be a signal for fear or pain, but may also be observed in lactating mothers protecting their nest. Determining the variation of the behavior from normalcy is a matter of training, studying, and observation.

TABLE A-1 An ethogram for *Macaca mulatta* (rhesus macaque)

Behavioral Categories	Recorded Behavior	De“nitions
Aggressive behaviors	Facial threat displays	Open mouth face ± bared teeth or vocalization
	Aggressive approach	Stiff approach, attacking run
	Physical aggression	Slap, grab, biting, or wrestling
Submissive behaviors	Facial submissive display	Bared teeth grin ± vocalization
	Avoidance	Avoid, ”ee, leave, displaced
	Active appeasement	Groom present, lip smacking
Af“liative behaviors	Af“liative contact	Contact sit (within arms reach), embrace, touch
	Passive grooming	Being groomed
	Active grooming	Grooming other animal
Sexual behaviors	Sexual contact	Genital present/inspection, mounting
Appetitive behaviors	Foraging	Food search, eating, drinking
Other activities	Active	Locomotion, enrichment use, self-grooming
Abnormal behaviors	Abnormal behaviors	Stereotypies, autoaggression
Inactive	Inactive	Lying, huddling, sitting, sleeping
Vigilance	Monitoring others	Visually following other individuals

Reprinted from Augustsson, A. and J. Hau. 1999. A simple ethological monitoring system to assess social stress in group-housed laboratory rhesus macaques. Lab Anim Sci 49:84-90.

Copyright © National Academy of Sciences. All rights reserved.

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

TABLE A-2 An ethogram for *Callithrix jacchus* (common marmoset)

Behavior	Code	De“nition
Agonism		
Tufts-”ick	TF	Rapid back-and-forth movement of ear tufts
Frown	FR	Lower eyebrows, furl brow, and turn down corners of mouth while staring
Cuff	CU	Swift, super“cial blow or scratch performed aggressively
Chase	CH	Pursue partner, withone or both animals exhibiting aggression and/or submission (not play)
Fight	FI	Grapple aggressively with partner(s), involving biting, clawing, and wrestling
Attack	AT	Lunge at or pounce on partner aggressively; may or may not result in “ght
Snap bite	SB	Direct a single short, sharp bite at partner
Submit	SU	Flatten ear tuftsand/or facial grimace (partially open mouth with corners of mouth retracted, exposing lower and sometimes upper teeth) and/or slit eyes (eyelids half closed)
Continuous submit FRE	CS	Continuous submit; start scoring after 5 sec

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

TABLE A-2 Continued

Behavior	Code	De“nition
Other social behaviors		
Sniff/nuzzle	SN	Orient face against or toward partner, excluding anogenital region
Anogenital inspect	AI	Orient face against ortoward anogenital region of partner, or use hands or mouthto investigate anogenital region of partner; includes anogenital groom
Groom	GR	Use hands and/or mouth topick through fur and/or mouth of partner, excluding anogenital region
Sexual solicit	SS	Stare at partner withear tufts ”attened and eyes slit
Mount	MO	Climb on partner•s backfrom behind and grip partner around waist and legs; may be accompanied by pelvic thrusting
Initiate huddle	IH	Establish passive, torso-torsbody contact with partner, with both animals remaining stationary andin passive contact for at least 3 sec
Leave huddle	LH	Terminate huddle after at least 3 sec of passive, torso-torso body contact during which both partners remained stationary
Object steal	OS	Take any nonfood objectfrom hands or mouth of partner
Attempt object steal	AO	Attempt but fail to take nonfood object from hands or mouth of partner
Food-associated behaviors		
Food steal	ST	Take any foodfrom hands or mouth of partner
Attempt food steal	AF	Attempt but fail to take food from hands or mouth of partner
Share food	SH	Eat from a food souce from which partner is simultaneously eating or that partner is occupying without removing any food from partner•s mouth or hands
New food	NF	Eat from a food souce that no other animal is currently holding, eating from, or occupying
Individual behaviors		
Bristle strut	BS	Arching posture and/or strut locomotion and/or general piloerection
Scent mark	SM	Rub or drag anogenital, suprapubic, or sternal region along substrate, object, or partner
Genital present	GP	Raise tail to expose genitals
Object manipulation	OM	Sniff, bite, chew, gouge,handle, pounce on, grapple with, or otherwise manipulate inanimate object, excluding food items and water bottle, for at least 1 sec

Written by Lissa Pabst. FromPrimate Info Net, Library and Information Service, National Primate Research Center, University of Wisconsin-Madison. Available at: http://pin.primate.wisc.edu/callicam/ethogram.html.

Copyright © National Academy of Sciences. All rights reserved.

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

TABLE A-3 An ethogram for rabbits

Common rabbit postures, behaviors, and vocalizations >	
Purring or teeth purring	„A sound made by lightly and

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

Other useful references for common normal behaviors of various labo< ratory animals include:

€ @Whishaw, I. Q. and B. Kolb, eds. 2005. The Behavior of the Labo= ratory Rat: A Handbook with Tests. New York: Oxford University Press.
€ @Pellis, S. M. and V. C. Pellis. 1987. Play-“ghting differs from serious “ghting in both target of attack and tactics of “ghting in the labora< tory rat Rattus norvegicus. Aggressive Behav 13:227-242.
€ @Bassett, L., H. M. Buchanan-Smith, and J. McKinley. 2003. Effects of training on stress-related behavior of the common marmoset (Callithrix jacchus) in relation to coping with routine husbandry procedures. J Appl Anim Welf Sci 6(3):221-233.
€ @670(Academico-7(Pon s.o-7(124069 64o-7(ppelude:)]TJ EMC /LI_Label

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

€ @Dixon, A. K. and H. U. Fisch. 1998. Animal models and ethological strategies for early drug testing in humans. Neurosci Biobehav Rev 23(2):345-358.

As stated in various places in the report, an effective assessment of distress is predicated upon solid knowledge of physiologic behaviors for each species and careful observation. In this respect, clinical and behavioral analysis of distress follows the investigative guidelines to determine the cause of any clinical symptomatology or pathology. Similarly, the goal of this exercise would be to remove, alleviate, or minimize the cause of distress (if doing so does not conflict with the research protocol) and support the animal in order to help it recover (see decision-making algorithm at the end of Chapter 4). The approach should integrate information from multiple behavioral and physiological parameters and should involve a team approach that includes researchers, veterinarians, and animal caretakers/technicians, as distress levels will vary in relation to the species, husbandry conditions, and experimental protocol as well as with each individual animal. The Committee points out that differential diagnosis of signs (clinical and behavioral) attributed to pain, sickness or distress is quite difficult and requires careful observation and clinical skills. The following tables showcase the overlapping clinical signs and abnormal behaviors associated both with distress and/or pain in various animal species.

Copyright © National Academy of Sciences. All rights reserved.

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

TABLE A-6 Common clinical signs associated with pain in small mammals

Production of fewer, smaller, or no fecal pellets	Reluctance to curl when sleeping (ferrets)
Anorexia	Tucked into abdomen

SHEET A-1 Investigational screen for behavioral phenotyping >

Behavioral Phenotyping Level 1 Screen *Accession #*_____

Date_____Investigator_____Genotype_____

Background strain(s) _____Inbred / N#___ Tg /TM KO/KI/Cond Gene Name _____

Key: 0 = zero; 1 = slow or reduced; ***2 = normal****; 3 = hyper*

Animal #		WT Hemi -/-	Animal #		WT Hemi -/-
DOB/Age		Sex M F	DOB/Age		Sex M F
Weight (g)		Fur color	Weight (g)		Fur color
Condition Score			Condition Score		
Empty Cage 2 mins:		Exploring 0..1..**2**..3	**Empty Cage 2 mins:**		Exploring 0..1..**2**..3
Gait abnormal	Y N	0= <1 side; 1 =< 1 circuit;	Gait abnormal	Y N	0= <1 side; 1 =< 1 circuit;
Posture abnormal	Y N	2= multiple circuits; 3= frantic	Posture abnormal	Y N	2= multiple circuits; 3= frantic
Freezing	Y N		Freezing	Y N	
Wild running	Y N	Digging 0..1..**2**..3	Wild running	Y N	Digging 0..1..**2**..3
Stereotypies	Y N	Grooming 0..1..**2**..3	Stereotypies	Y N	Grooming 0..1..**2**..3
Escape	Y N	Rearing 0..1..**2**..3	Escape	Y N	Rearing 0..1..**2**..3

DORSAL VENTRAL DORSAL VENTRAL



DRAW DRAW

Bald patches/abnorm alities Bald patches/abnorm alities

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

The following score sheets have been developed to assess animals in toxicology studies. This assessment is based on a detailed and systematic observation scheme that identifies and scores abnormalities according to

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

SHEET A-2 Continued

l. @ Abnormal skin or hair-coat/mucous membranes: Description of atypical skin or mucous membrane color, changes in hair coat, loss of fur, etc.
m. @ Excessive soiling: Description and location of increased body soiling.
n. @ General abnormalities: Description of any other atypical “nding not “tting any of the previous observation categories.

Step 2. Detailed Clinical Observations (DCO)

The purpose of the DCO examination is to provide information on the physical health of the animals for the duration of a study, as well as to document any changes in health status that may have occurred in response to chemical treatment of the animals. This examination, scheduled periodically during a study, is conducted in a careful and systematic manner. The examination begins at the head of the animal and works toward the rear of the animal. The observations are ranked according to severity.

A. = Cage-side observations.
€ @ Abnormal movements or behaviors: Unusual body movements (e.g., tremors, convulsions), abnormal behaviors (e.g., circling, stereotypy) and changes in posture (e.g., arched back, splayed stance).
€ @ Resistance to removal: The degree to which the animal attempts to escape capture is scored. The observer will slowly present a gloved hand into the cage and will grasp the animal over the shoulder area or by the tail.
1 = Decrease„clearly less resistance to capture than typical
2 = Typical„minimally to actively avoids capture and may be mildly aggressive
3 = Increase„clearly more resistance to capture than typical and is very aggressive (attempts to bite)
B. = Hand-held observations recorded while handling an animal.
1. @ Ranked observations„the following use a de“ned scale to rank the degree of severity:
a. > Eye observations Eyes are bilaterally examined; however, if a unilateral observation is made, a concurrent observation is not made for the other eye if it is within typical limits.
(1) @ Palpebral closure >
1 = Closed (50% to completely closed) >
2 = Open >
3 = Protruding eyes >
(2) @ Pupil size (aided by penlight): Under typical examination conditions (white light), the typical appearance of the pupils in albino animals is complete constriction. Therefore a decrease in pupil size cannot be observed.
0 = Unable to evaluate
1 = Decrease„clearly decreased pupil size compared to typical
2 = Typical„completely constricted pupils
3 = Increase„clearly increased pupil size compared to typical
(3) @ Lacrimation (noncolored periocular wetness)
1 = Decrease„extremely dry appearance of cornea
2 = Typical„glistening cornea (moderate dryness or wetness)
3 = Increase„extensive wetness around the eyes

continued

Copyright © National Academy of Sciences. All rights reserved.

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

An alternative example is the following score sheet developed at the University of Birmingham Biomedical Sciences Unit, courtesy of David B. Morton, BVSc, PhD.

SHEET A-3 General screening and applicability 986 Tm [(Str)30(ain: Start)]TJ 24 1

Isolated							

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

SHEET A-3 Continued

Hunched posture							
Boarding of abdomen							
Tiptoe walking							
Handling							
Aggressive							
Not inquisitive/alert							
Crusty eyes/nose							
Reluctant to move							
Dehydration/skin tone							
Vocalisation							
Hyperactive							
Bodyweight							
% weight change from start							
% weight change previous day							
Body temperature							
Treatment							
Prolapse							
Overgrown teeth							
Other							
NAD							
Vet/PI contacted							
Signature:							

Scoring details
* Abnormal breathing: Breaths/min
** Type of breathing: R: Rapid,S: Shallow, L: Labored, N: Normal

Copyright © National Academy of Sciences. All rights reserved.

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

As stated in Chapter 4, establishing surrogate or humane endpoints as part of the experimental protocol andbefore experiments commence is one



RecogMCID 0 >

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

genetic backgrounds and disease states. She is particularly interested in the role of genetics on complex behavioral phenotypes and has been working with mouse models of human disease for over 16 years. She has previously served as a member of the research ethics committee at St. Mary's University, Nova Scotia, Canada.

Marilyn J. Brown, DVM, is Executive Director of Animal Welfare and Training at Charles River Laboratories. She is a Diplomate of the American College of Laboratory Animal Medicine. She has experience dealing with issues pertaining to animal welfare, as well as serving on a number of NRC committees, including the ILAR Committee to Revise the Guide for the Care and Use of Laboratory Animals (1996). She also served as a reviewer for the 1992 ILAR report Recognition and Alleviation of Pain and Distress in Laboratory Animals. Dr. Brown has been the recipient of numerous honors and awards, including being awarded the title of De Facto Diplomate for the European College of Laboratory Animal Medicine.

Fon Chang DVM, is Senior Staff Veterinarian at AstraZeneca Pharmaceuticals LP. He completed residencies in Laboratory Animal Medicine and in Animal Behavior at UC-Davis. His experience on animal welfare and behavior issues includes behavioral management of research animals and behavioral phenotyping of genetically engineered mice. Dr. Chang's professional affiliations include the American Association for Laboratory Animal Science, the American Society of Laboratory Animal Practitioners, and the American Veterinary Medical Association.

James P. Herman, PhD, is Professor of Psychiatry and Director of the Laboratory of Stress Neurobiology at the University of Cincinnati. His research pertains to rodents and the structural and functional neurocircuitry involved in the stress response, molecular effects of stress in the brain, and gender differences in the stress response. Dr. Herman was recently a panel member of the American Physiological Society Workshop on Managing Pain and Distress in Experimental Animals, and has been an invited speaker on the topic of distress for the AALAS and PRIM&R/ARENA national meetings.

Robert Hubrecht, PhD, has been with the Universities Federation for Animal Welfare (UFAW) since 1992, where he has served as Deputy Director since 1996 (UFAW is a unique scientific and technical animal welfare organization in the UK, which uses scientific knowledge to improve the welfare of animals kept as pets, in zoos and laboratories, and on farms). He has a long career in the fields of animal welfare and ethology. This includes, in addition to a number of academic positions, service as a member of the Animal Procedures Committee (APC). The APC advises the UK Home

Copyright © National Academy of Sciences. All rights reserved.

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

Index >

A

Abnormal behavior. See also speci“c behaviors
arousal reduction hypothesis, 35 >
clinical signs of distress, 34-35, 103 >
examples, 32-33 >
habit hypothesis, 35 >
observation strategies, 33, 89-90 >
recognizing, 4, 11, 19, 33-36 >
research on, 4, 6, 36, 88 >
as signs of distress, 3, 4, 16, 19, 35, 103
as signs of stress, 2, 3-4, 14, 16, 36 >
species-speci“c, 103 >
stimulation hypothesis, 35 >
Adrenocorticotropic hormone (ACTH), 29 > n.1, 36-37 >
Age, and distress response, 3, 4, 6, 25, 27, > 33, 34, 35, 41, 43, 88, 95-96, 105 >
Aggression, 3, 4, 16, 26, 28, 30, 68-69, 96, > 97, 99, 103, 104, 107, 108, 110, > 112 >
Alleviating distress.See alsoEuthanasia >
decision and response algorithm, 43, 75, > 76 >
experimental design and, 70-71 >
recommendations, 6 >
removal of animal from study, 5, 19, 43, > 75 >
team approach, 75-78 >
treatment modalities, 66, 75-77 >
American Association for Laboratory Animal > Science, 8, 92 >
American College of Laboratory Animal > Medicine, 8, 92 >
Analgesics, 70, 75 >
Animal training, 6, 75, 76, 90 >
Animal welfare >
de“ning, 17-21 >
descriptors, 18-19 >
distress and, 17-21, 33, 88 >
historical and cultural context, 1, 17-19 > > institutional support, 7, 91 >
Animal Welfare Act, 1, 10, 13 >
Animal Welfare Institute, 7, 8, 91, 92 >
Anthropocentrism, 90 >
Anthropomorphism, 20, 90 >
Anxiety-related behavior, 21, 28, 30, 66-67, > 79 >
Anxiolytics, 66, 75, 77 >
Arousal reduction hypothesis, 35 >
Assessment of stress and distress >
behavioral, 19, 26, 95-100 >
considerations in, 4, 18 >
de“ning distress and, 3, 18, 88 >
experimental design and, 4, 43 >
information resources, 100-101 >
physiological measures, 19-20 >
process, 43-44, 106-108 >
sampling procedures, 36, 37-38, 39, 60 >
team approach, 4, 42, 43, 44 >
tools for, 88, 95-112 >

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

Association for the Assessment and > Accreditation of Laboratory Animal > Care International, 1, 10 >
Autism, 35 >
Autonomic nervous system, 4, 28, 39, 40, 41>
Avoidance/escape behavior, 20, 36, 96, 99, > 105, 107 >
Avoiding/minimizing distress >
in animal care, 5, 64-70 >
in animal use, 5, 70-75 >
in clinical sampling procedures, 36, 37< 38, 39, 43, 70-71 >
enrichment, 5, 6, 7, 11, 19, 31, 32-33, > 35, 63, 64, 66-67, 69, 75, 77, 90, 91, > 96 >
in experimental design, 5, 6, 11-12, 22, > 63, 70-74, 90 >
housing, 11, 18, 19, 43, 63, 65-66, 90 >
humane endpoints, 5, 11-12, 34, 63, 71, > 88, 111 >
husbandry, 5, 6, 18, 19, 63, 64, 68, 69< 70, 75, 90 >
interference with research aims, 63, 64, > 67 >
minimally or noninvasive technologies, > 5, 39, 74-75 >
pilot studies, 5, 67, 72 >
regulatory requirements, 1, 2, 5, 9-11, > 12, 75, 77 >
research, 64 >
sample size determination, 63, 72-74 >
socialization, 5, 11, 19, 63, 68-69, 90 >
statistical analyses and, 5, 7, 12, 63, 72< 74, 91 >
team approach, 5, 63 >
Three Rs, 5, 6, 63, 64, 89, 90 >

B

Bar chewing, 31, 32 >
Bar circling, 31 >
Barbering, excessive, 32 >
Behavior. SeeAbnormal behavior; Normal > behavior; speci“c behaviors <
assessment, 20 >
phenotyping, 104-105 >
Brain >
function, 19 n.4, 21, 29, 31-32, 41 >
physiological response to stress, 39-40, > 41, 87 >
size and structure, 28, 52 >
Brambell Committee, 18 >
Buprenorphine, 77 >

C

Cage cleaning, 69 >
Caloric restriction protocols, 34-35 >
Cardiovascular disease models, 75 >
Cats, 68, 102, 103 >
Chlorpromazine, 77 >
Coat/hair/fur >
appearance, 4, 27, 28, 34, 105, 107, > 109 >
sampling, 37, 38, 56, 70, 71 >
Cognitive de“cits, 21, 27, 28, 52 >
Colitis, 39, 61 >
Clinical signs of distress.See alsoPhysiology >
alleviating, 77 >
assessment tools, 102, 104 >
behavioral, 32-33, 34-35 >
identifying and interpreting, 43, 87-88, > 95, 104 >
sampling procedures, 36, 37-38, 39, 43, > 70-71 >
species-speci“c, 102 >
subclinical, 3, 16 >
Coprophagy70-71 >Coprocococococococococo79c022 Tw T*

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

Humane endpoints. See alsoEuthanasia >
and distress predictors, 88 >
and mice, 71, 111 >
avoiding/minimizing distress, 5, 11-12, > 34, 63, 71,88,111 >
clinical signs, 34 >
determining/establishing, 71, 88, 111 >
primates, 34 >
statement of task, 2, 10 >
tiered scoring system, 111 >
Husbandry, 4, 11, 43 >
avoiding/minimizing distress, 5, 6, 18, > 19, 63, 64, 68, 69-70, 75, 90 >
genetically manipulated animals, 89 >
prior to euthanasia, 77-78 >
research needs, 89-90 >
5-Hydroxyindoleacetic acid, 30 >
Hypertension, 3, 16-17 >
Hypothalamic-pituitary-adrenal (HPA) axis > activity, 16, 20-21, 27, 29 n.1, 31, > 36-37, 38, 40, 68 >

I ;

Immunological changes, 3, 4, 14, 16-17, 21, > 31, 34, 37, 41-42, 67 >
Impulse control, impaired, 30 >
In”ammatory disease models, 75 >
Institutional Animal Care and Use > Committees, 2, 5, 10, 12, 43, 63, 64, > 75, 79, 89 >
Interleukin-1, 42 >

J

Johns Hopkins Center for Alternatives to > Animal Testing, 8, 92 >

L

Learning abilities, 28, 34, 43, 52,53, 54, 70 >
Leukemia models, 71 >
Light conditions, 65, 78 >
Locomotor activity, 26, 27, 28, 31, 34, 35, > 52 >
Luteinizing hormone, 38, 58 >

M

Macaques, 30, 34, 68, 75, 96 >
Marmosets, 68, 96, 97-98, 100 >
Maternal behavior/environment, 25, 26, > 29-31, 33, 37 >
A-Melanocyte stimulating hormone [A< MSH], 38, 58 >
Memory, 21, 28 >
3-Methoxy-4-hydroxyphenylglycol, 29 n.1 >
Mice >
behavioral assessment, 88-89, 104, > 105-106 >
behavioral signs of distress, 26, 32, 66, > 77, 103 >
clinical signs of distress, 102 >
collagen-induced arthritis model, 111 >
cross-fostered, 29-30 >
enrichment, 32, 66-67 >
housing, 31, 66, 68, 69 >
humane endpoints, 71, 111 >
Huntington•s disease model, 26, 31 >
husbandry, 89-90 >
hyperthermia model, 28 >
inbred strains, 15, 26, 28, 73, 104 >
intraspecies differences in response, 15, > 26, 27, 28 >
normal behavior, 26, 27, 29, 31, 32 >
scoring sheets, 109-110, 111 >
socialization, 68, 69 >
transgenic and knockout models, 28, 29, > 31, 66-67, 88-89, 105 >

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

Neurotransmitters, 39-41 >
Noise stressors, 65-66, 78 >
Normal behavior >
age and, 4, 27, 34, 52, 95, 97 >
de“ned, 65 >
ethograms, 26, 52, 95-100 >
gender and, 4, 26, 27, 33, 34, 95 >
genetic variation, 4, 25, 26, 27, 28-29, > 34, 52, 95 >
housing and, 26, 31-33 >
infant-related, 97 >
identifying, 4, 11, 25, 26-33, 43 >
physiological state and, 4, 26, 31, 33, 34 >
rearing and postnatal separation, 26, > 29-31 >
tools for assessing, 96-112 >
transgenic and knockout models, 4, 25, > 26, 29 >

O

Oxytocin, 38, 58 >

P

Pain, 21 >
addressed separately from distress, 1, 9, > 10, 11 >
assessment, 102-104 >
chronic neuropathic, 38 >
clinical signs, 104 >
markers, 38 >
sensitivity, 18, 27, 28, 52 >
Peer rearing, 30-31 >
Pharmacological interventions, 66, 75, 77 >
Physiology. See alsoBrain; Clinical signs of > distress; Neurological changes >
aging and, 27 >
and behavior, 4, 26, 27, 31, 33 >
endocrinological changes, 28-29, 34, > 36-38 >
immunological changes, 3, 4, 14, 16-17, > 21, 31, 34, 37, 41-42, 67 >
measures of stress and distress, 2, 3, 4, > 16, 18, 19-21, 36-42, 56 >
subclinical changes, 16-17 >
telemetric monitoring, 39, 61 >
Pigs, 31, 66, 68 >
Pilot studies, 5, 67, 72 >
Play behavior, 27, 28, 97, 100 >
Postnatal separation.SeeRearing >
Poultry, 66 >
Prepulse inhibition behavior, 31-32 >
Preventing distressSeeAlleviating distress; > Avoiding/minimizing distress >
Primate Enrichment Database, 7, 91 >
Primates, nonhuman.See alsoMacaques; > Marmosets; Rhesus monkeys; Vervet > monkeys >
alleviating distress in, 76, 77 >
enrichment, 31 >
ethograms, 100, 102-103 >
high-reactor monkeys, 28-29 >
humane endpoints, 34 >
normal vs. abnormal behavior, 25, 26, > 28-29, 30-31, 32, 96, 112 >
pair testing record, 112 >
rearing environment, 30-31 >
sample collection methods, 70-71 >
Prolactin, 2, 4, 14, 38, 58-60 >

R

Rabbits, 31, 66, 68, 96,99, 100, 102, 103, > 104 >
Rats >
clinical signs of distress, 102-103 >
cross-fostered, 29-30 >
habituation to test environment, 21 >
housing, 31, 32, 68 >
husbandry, 69 >
normal behavior, 27, 30, 31, 32 >
physiologic effects of stress, 27, 40, 41 >
shock stimuli response, 16, 40, 41 >
socialization, 68 >
Rearing and postnatal separation, 25, 29-31 >
Recognition of stress and distress >
behavioral correlates, 3-4, 11, 19, 26-36; > see alsoAbnormal behavior; Normal > behavior >
Recommendations, 6-8, 90-92 >
Regulatory requirements, 1, 2, 9-11, 75, 77 >
Regurgitation/rumination, 32 >

Recognition and Alleviation of Distress in Laboratory Animals
http://www.nap.edu/catalog/11931.html

publication of, 7, 91
scope of this study, 1-2, 10-11
Restraint of animals, 15, 16, 19, 37, 38, 41<
42, 69, 70, 108
Rhesus monkeys, 25, 28-29 n.1, 32, 75, 77,
96, 100

S

S6 kinase gene, 29
Sample size determination, 22, 63, 64, 72<
75, 90
Schizophrenia, 35
Selective serotonin reuptake inhibitors, 77
Self-injurious behavior, 4, 32, 35, 75, 77, 95